产品货号: AMM00731



产品概述 (Summary)

产品名称 (Production Name) CD4 (7H9) Mouse Monoclonal Antibody

描述 (Description) Mouse monoclonal Antibody

宿主 (Host) Mouse 应用 (Application) IHC

种属反应性 (Reactivity) Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) lgG1

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

sodium azide, pH 7.3.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CD4

CD4; T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; CD 别名 (Alternative Names)

antigen CD4

基因 ID (Gene ID) 920 **蛋白 ID (SwissProt ID)** P01730.

产品应用(Application)

稀释比 (Dilution Ratio) IHC 1:50-1:100

蛋白分子量(Molecular Weight) -

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品货号: AMM00731




Cluster of Differentiation 4 (CD4) is a glycoprotein composed of an amino-terminal extracellular domain (four domains: D1-D4 with Ig-like structures), a transmembrane part and a short cytoplasmic tail. CD4 is expressed on the surface of T helper cells, regulatory T cells, monocytes, macrophages and dendritic cells, and plays an important role in the development and activation of T cells. On T cells, CD4 is the co-receptor for the T cell receptor (TCR), and these two distinct structures recognize the Antigen–Major Histocompatibility Complex (MHC).

研究领域(Research Area)

Immunology

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded Human stomach tissue using CD4 (7H9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Human tonsils using CD4 (7H9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品货号: AMM00731






Immunohistochemistry analysis of paraffin-embedded mouse brain tissue using CD4 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunofluorescence analysis of CD4 (7H9) in mouse colon tissue using CD4 (7H9) antibody(11A1)(red), and DAPI (blue).




Immunofluorescence analysis of CD4 (7H9) in rat lung using CD4 antibody(11A1)(red), and DAPI (blue).

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品货号: AMM00731





Immunohistochemistry analysis of paraffin-embedded Human Amygdala using CD4 (7H9) antibody. High-pressure and temperature Tris-EDTA pH 8.0 was used for antigen retrieval.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838