

产品名称: Hsc70 (4C2) Mouse Monoclonal Antibody
产品货号: AMM00897



产品概述 (Summary)

产品名称 (Production Name)	Hsc70 (4C2) Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	HSPA8
别名 (Alternative Names)	Heat shock cognate 71 kDa protein (Heat shock 70 kDa protein 8)
基因 ID (Gene ID)	3312
蛋白 ID (SwissProt ID)	P11142.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000,IHC 1:50-1:100
蛋白分子量 (Molecular Weight)	Calculated MW: 71 kDa; Observed MW: 71 kDa

研究背景 (Background)

Acts as a repressor of transcriptional activation. Inhibits the transcriptional coactivator activity of CITED1 on Smad-

产品名称: Hsc70 (4C2) Mouse Monoclonal Antibody
产品货号: AMM00897




mediated transcription. Chaperone. Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing.


研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded Human Breast Carcinoma using Hsc70 (4C2) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of Hsc70 (4C2) in HeLa, A549, HL-60, U2OS, C6 lysates using Hsc70 (4C2) antibody

注意事项 (Note)

For research use only .