

产品名称: LC3A/B (3E9) Mouse Monoclonal Antibody
产品货号: AMM03560



产品概述 (Summary)

产品名称 (Production Name)	LC3A/B (3E9) Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG2b
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MAP1LC3A/MAP1LC3B
别名 (Alternative Names)	LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A; LC3B; ATG8F; MAP1LC3B-a; MAP1A/1BLC3; MAP1LC3B
基因 ID (Gene ID)	84557/81631
蛋白 ID (SwissProt ID)	Q9H492/Q9GZQ8.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000
蛋白分子量 (Molecular Weight)	Calculated MW: 14 kDa; Observed MW: 14,16 kDa

研究背景 (Background)

产品名称: LC3A/B (3E9) Mouse Monoclonal Antibody
产品货号: AMM03560




Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.

研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Western blot analysis of LC3A/B in rat Cerebral Cortex, rat Brain and Hela lysates using LC3A/B antibody.

注意事项 (Note)

For research use only .