

产品名称: Ubiquitin(5F1)Mouse Monoclonal Antibody
产品货号: AMM19553



产品概述 (Summary)

产品名称 (Production Name)	Ubiquitin(5F1)Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ICC/IF
种属反应性 (Reactivity)	Human,Rat,Mouse,Other

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	
别名 (Alternative Names)	
基因 ID (Gene ID)	7314.0
蛋白 ID (SwissProt ID)	PAN.Synthetic Peptide of Ubiquitin

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:100-1:200
蛋白分子量 (Molecular Weight)	

研究背景 (Background)

UBB (ubiquitin B) encodes ubiquitin, one of the most conserved proteins known. Ubiquitin has a major role in targeting


产品名称: Ubiquitin(5F1)Mouse Monoclonal Antibody
产品货号: AMM19553




cellular proteins for degradation by the 26S proteasome. It is also involved in the maintenance of chromatin structure, the regulation of gene expression, and the stress response. Ubiquitin is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin moiety fused to an unrelated protein. UBB consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. An aberrant form of this protein has been detected in patients with Alzheimer's disease and Down syndrome. Pseudogenes of UBB are located on chromosomes 1, 2, 13, and 17. Alternative splicing results in multiple transcript variants.

研究领域 (Research Area)

图片 (Image Data)




Immunohistochemical analysis of paraffin-embedded Human-Tonsil tissue. 1, Ubiquitin Mouse Monoclonal Antibody (5F1) was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, Secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1, Ubiquitin Mouse Monoclonal Antibody (5F1) was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, Secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.


产品名称: Ubiquitin(5F1)Mouse Monoclonal Antibody
产品货号: AMM19553



Immunofluorescence analysis of Human-stomach-cancer tissue. 1,Ubiquitin Mouse Monoclonal Antibody (5F1) (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of Mouse-brain tissue. 1,Ubiquitin Mouse Monoclonal Antibody (5F1) (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of Rat-brain tissue. 1,Ubiquitin Mouse Monoclonal Antibody (5F1) (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B


产品名称: Ubiquitin(5F1)Mouse Monoclonal Antibody
产品货号: AMM19553



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma Tissue using Ubiquitin Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Human Stomach Carcinoma Tissue using Ubiquitin Mouse mAb diluted at 1:200.



Western blot analysis of 1) Hela Cell Lysate, 2) 3T3 Cell Lysate, 3) Rat Brain Tissue Lysate using Ubiquitin Mouse mAb diluted at 1:1000.

注意事项 (Note)

For research use only .