产品名称: FGF2 Mouse Monoclonal Antibody

产品货号: AMM80535



产品概述 (Summary)

产品名称 (Production Name) FGF2 Mouse Monoclonal Antibody

描述 (Description) Mouse monoclonal Antibody

宿主 (Host)Mouse应用 (Application)IHC,ELISA种属反应性 (Reactivity)Human

产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG1 充隆 (Clonality) Monoclonal Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

储存溶液 (Buffer) Purified antibody in PBS with 0.05% sodium azide.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

存放说明 (Storage)

基因名 (Gene Name) FGF2

别名 (Alternative Names) BFGF; FGFB; HBGF-2; FGF2

基因 ID (Gene ID) 2247.0

蛋白ID (SwissProt ID) P09038.Purified recombinant fragment of FGF2 expressed in E. Coli.

产品应用(Application)

稀释比 (Dilution Ratio) IHC 1:200-1:1000,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) /

研究背景 (Background)

FGF2 is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. FGF2 is a single-chain polypeptide growth factor that plays a significant role in the

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: FGF2 Mouse Monoclonal Antibody

产品货号: AMM80535




process of wound healing and is a potent inducer of anguogenesis. Due to its basic pH, the factor is named FGF-2 (basic FGF, bFGF). Several different forms of the human protein exist ranging from 18-24 kDa in size due to the use of alternative start sites within the fgf-2 gene. It has a 55 percent amino acid residue identity to FIBROBLAST GROWTH FACTOR 1 and has potent heparin-binding activity. The growth factor is an extremely potent inducer of DNA synthesis in a variety of cell types from mesoderm and neuroectoderm lineages. It was originally named basic fibroblast growth factor based upon its chemical properties and to distinguish it from acidic fibroblast growth factor (FIBROBLAST GROWTH FACTOR 1).

研究领域 (Research Area)

TGF-beta signaling pathway, PI3K-Akt signaling pathway, MAPK signaling pathway

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human recturn adenocarcinoma tissue showing cytoplasmic localization using FGF2 mouse mAb with DAB staining.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838