

产品名称: Vimentin Mouse Monoclonal Antibody
产品货号: AMM80578



产品概述 (Summary)

产品名称 (Production Name)	Vimentin Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ELISA,FC
种属反应性 (Reactivity)	Human,Mouse,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS containing 0.03% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	Vimentin
别名 (Alternative Names)	FLJ36605; VIM
基因 ID (Gene ID)	7431.0
蛋白 ID (SwissProt ID)	P08670.Purified recombinant fragment of Vimentin expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	54kDa

研究背景 (Background)

Vimentin is the major subunit protein of the intermediate filaments of mesenchymal cells. It is believed to be involved with the intracellular transport of proteins between the nucleus and plasma membrane. Vimentin has been implicated to be


产品名称: **Vimentin Mouse Monoclonal Antibody**
产品货号: **AMM80578**




involved in the rate of steroid synthesis via its role as a storage network for steroidogenic cholesterol containing lipid droplets. Vimentin phosphorylation by a protein kinase causes the breakdown of intermediate filaments and activation of an ATP and myosin light chain dependent contractile event. This results in cytoskeletal changes that facilitate the interaction of the lipid droplets within mitochondria, and subsequent transport of cholesterol to the organelles leading to an increase in steroid synthesis. Immunohistochemical staining for Vimentin is characteristic of sarcomas (of neural, muscle and fibroblast origin) compared to carcinomas which are generally negative. Melanomas, lymphomas and vascular tumors may all stain for Vimentin. Vimentin antibodies are thus of value in the differential diagnosis of undifferentiated neoplasms and malignant tumors. They are generally used with a panel of other antibodies including those recognising cytokeratins, lymphoid markers, S100, desmin and neurofilaments.

研究领域 (Research Area)

图片 (Image Data)




Western blot analysis using Vimentin mouse mAb against NIH/3T3 cell lysate.



Flow cytometric analysis of COS7 cells using Vimentin mouse mAb (green) and negative control (red).

产品名称: Vimentin Mouse Monoclonal Antibody
产品货号: AMM80578



Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue, showing cytoplasmic localization using Vimentin mouse mAb with DAB staining.

注意事项 (Note)

For research use only .