

产品名称: IGF1R-Beta Mouse Monoclonal Antibody
产品货号: AMM80596



产品概述 (Summary)

产品名称 (Production Name)	IGF1R-Beta Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	IHC,ELISA
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG2b
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS containing 0.03% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IGF1R-Beta
别名 (Alternative Names)	IGF1R, IGF1R-Beta
基因 ID (Gene ID)	3480.0
蛋白 ID (SwissProt ID)	P08069.Purified recombinant fragment of IGF1R-Beta expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	96kDa

研究背景 (Background)

IGF1R (insulin-like growth factor 1 receptor), a transmembrane receptor tyrosine kinase, is widely expressed in many cell types within fetal and postnatal tissues, and in many cell lines. Upon binding to its ligands, IGF-I and IGF-II, receptor

产品名称: IGF1R-Beta Mouse Monoclonal Antibody
产品货号: AMM80596




autophosphorylation occurs. The triple tyrosine cluster within the kinase domain (Tyr1131, Tyr1135 and Tyr1136) is the earliest major site of autophosphorylation. Phosphorylation of these three tyrosine residues is necessary for kinase activation. Insulin receptors (IRs) share significant similarity with IGF1 receptors in both structure and function, including an equivalent triple tyrosine cluster within the activation loop of the kinase domain (Tyr1146, Tyr1150 and Tyr1151). Tyrosine autophosphorylation of insulin receptor is one of the earliest cellular responses to insulin stimulation. Autophosphorylation begins with phosphorylation of Tyr1146 and either Tyr1150 or Tyr1151. Full kinase activation requires the triple tyrosine phosphorylation.

研究领域 (Research Area)

PI3K-Akt signaling pathway, Jak-STAT signaling pathway, Hippo signaling pathway

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human gastric adenocarcinoma(A), colon adenocarcinoma(B), endometrial carcinoma(uterus)(C), ovary adenocarcinoma(D), lung squamous cell carcinoma(E), stomach epithelium mucosae(F), showing membrane localization using IGF1R-Beta mouse mAb with DAB staining.

注意事项 (Note)

For research use only .