

产品名称: TIP60 Mouse Monoclonal Antibody
产品货号: AMM80598



产品概述 (Summary)

产品名称 (Production Name)	TIP60 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	IHC,ELISA
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG2b
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS containing 0.03% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TIP60
别名 (Alternative Names)	TIP60 (HTATIP)
基因 ID (Gene ID)	10524.0
蛋白 ID (SwissProt ID)	Q92993.Purified recombinant fragment of human TIP60 expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	60kDa

研究背景 (Background)

HTATIP (HIV-1 Tat interacting protein TIP60, about 60kDa) belongs to the MYST family of histone acetyl transferases (HATs) and was originally isolated as an HIV-1 TAT-interactive protein. HATs play important roles in regulating chromatin


产品名称: TIP60 Mouse Monoclonal Antibody
产品货号: AMM80598



remodeling, transcription and other nuclear processes by acetylating histone and nonhistone proteins. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. In addition to the growing number of post-translational histone modifications regulating chromatin structure, cells can also exchange canonical histones with variant histones that can directly or indirectly modulate chromatin structure. There are five major variants of histone H2A: canonical H2A (most abundant), H2A.X, MacroH2A, H2ABbd and H2A.Z. Histone H2A.Z, the most conserved variant across species, functions as both a positive and negative regulator of transcription and is important for chromosome stability. Several homologous protein complexes, such as SWR-C, TIP60 and SRCAP (mammals), have been shown to catalyze the ATP-dependent exchange of H2A.Z for H2A in the nucleosome. This protein is a histone acetylase that has a role in DNA repair and apoptosis and is thought to play an important role in signal transduction.

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human liver carcinoma (A), rectum carcinoma (B), normal medulla tissue (C) and normal interbrain tissues (D), showing nuclear localization using Tip60 mouse mAb with DAB staining.

注意事项 (Note)

For research use only .