产品名称: ITGA5 Mouse Monoclonal Antibody

产品货号: AMM80787



产品概述 (Summary)

产品名称 (Production Name) **ITGA5 Mouse Monoclonal Antibody**

描述 (Description) Mouse monoclonal Antibody

宿主 (Host) Mouse

应用 (Application) IHC,ELISA,FC

种属反应性 (Reactivity) Human

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG2a Monoclonal 克隆 (Clonality) 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

储存溶液 (Buffer) Purified antibody in PBS with 0.05% sodium azide.

纯化方式 (Purification) **Affinity Purification**

免疫原信息 (Immunogen)

存放说明 (Storage)

基因名 (Gene Name) ITGA5

别名 (Alternative Names) FNRA; CD49e; VLA5A; ITGA5

基因 ID (Gene ID) 3678.0

蛋白 ID (SwissProt ID) P08648.Purified recombinant fragment of human ITGA5 expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio) IHC 1:200-1:1000, ELISA 1:5000-1:20000, FC 1:200-1:400

蛋白分子量 (Molecular Weight) 114kDa

研究背景 (Background)

The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This gene encodes the integrin alpha 5 chain. Alpha chain 5

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838 产品名称: ITGA5 Mouse Monoclonal Antibody

产品货号: AMM80787




undergoes post-translational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains that join with beta 1 to form a fibronectin receptor. In addition to adhesion, integrins are known to participate in cell-surface mediated signalling. Integrin alpha 5 is a heterodimer that associates noncovalently with CD29/integrin beta 1 subunit to form the alpha-5-beta-1 very late antigen (VLA-5) complex. VLA-5 is a fibronectin receptor that is expressed on thymocytes, T-cells, monocytes and platelets. It is also found on very early B-cells and activated B-cells. VLA-5-mediated binding to fibronectin sends a costimulatory signal to T-cells and enhances Fc-gamma-R- and complement receptor-mediated phago-cytosis. It is also involved in monocyte migration into extracellular tissues.


研究领域 (Research Area)

PI3K-Akt signaling pathway

图片 (Image Data)



Flow cytometric analysis of Hela cells using ITGA5 mouse mAb (green) and negative control (purple).



Immunohistochemical analysis of paraffin-embedded human Placenta tissues using ITGA5 mouse mAb

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838