产品名称: CDC2 Mouse Monoclonal Antibody

产品货号: AMM80840



产品概述 (Summary)

产品名称 (Production Name) CDC2 Mouse Monoclonal Antibody

描述 (Description) Mouse monoclonal Antibody

宿主 (Host) Mouse

应用 (Application) WB,IHC,ELISA

种属反应性 (Reactivity) Human

产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG1 充隆 (Clonality) Monoclonal Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) PBS containing 0.03% sodium azide.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CDC2

别名 (Alternative Names) CDC2; CDC28A; P34CDC2; MGC111195; DKFZp686L20222; CDK1

基因 ID (Gene ID) 983.0

蛋白ID (SwissProt ID) P06493.Purified recombinant fragment of CDC2 expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:200-1:1000,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 34kDa

研究背景 (Background)

The cell division control protein cdc2, also known as cyclin-dependent kinase 1 (Cdk1) or p34/cdk1, plays a key role in the control of the eukaryotic cell cycle, where it is required for entry into S-phase and mitosis. Cdc2 exists as a complex with

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CDC2 Mouse Monoclonal Antibody

产品货号: AMM80840




both cyclin A and cyclin B. The best characterized of these associations is the Cdc2 p34 cyclin B complex, which is required for the G2 to M phase transition. Activation of Cdc2 is controlled at several steps including cyclin binding and phosphorylation of threonine 161. However, the critical regulatory step in activating cdc2 during progression into mitosis appears to be dephosphorylation of Tyr15 and Tyr14. Phosphorylation at Tyr15 and inhibition of Cdc2 is carried out by WEE1 and MIK protein kinases while Tyr15 dephosphorylation and activation of Cdc2 is carried out by the cdc25 phosphatase. The isoform CDC2deltaT is found in breast cancer tissues. Furthermore, cdc2/Cdk1 is a key mediator of neuronal cell death in brain development and degeneration.


研究领域 (Research Area)

Apoptosis

图片 (Image Data)



Western blot analysis using CDC2 mouse mAb against Jurkat (1) cell lysate.



Immunohistochemical analysis of paraffin-embedded human lung cancer (A) and esophageal cancer (B), showing cytoplasmic localization using CDC2 mouse mAb with DAB staining.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838