

产品名称: CDKN2A Mouse Monoclonal Antibody
产品货号: AMM81317



产品概述 (Summary)

产品名称 (Production Name)	CDKN2A Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ELISA,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CDKN2A
别名 (Alternative Names)	ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A
基因 ID (Gene ID)	1029.0
蛋白 ID (SwissProt ID)	P42771.Purified recombinant fragment of human CDKN2A (AA: 1-156) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	16.5kDa

研究背景 (Background)


产品名称: CDKN2A Mouse Monoclonal Antibody
产品货号: AMM81317




This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, MDM1, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

研究领域 (Research Area)

图片 (Image Data)




Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);




Western blot analysis using CDKN2A mouse mAb against Hela (1), HepG2 (2) and Hek293 (3) cell lysate.


产品名称: CDKN2A Mouse Monoclonal Antibody
产品货号: AMM81317



Flow cytometric analysis of Hela cells using CDKN2A mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human endometrial cancer tissues using CDKN2A mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human esophageal cancer tissues using CDKN2A mouse mAb with DAB staining.

注意事项 (Note)

For research use only .