

产品名称: PRKACA Mouse Monoclonal Antibody
产品货号: AMM81371



产品概述 (Summary)

产品名称 (Production Name)	PRKACA Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	ELISA
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PRKACA
别名 (Alternative Names)	PKACA
基因 ID (Gene ID)	5566.0
蛋白 ID (SwissProt ID)	P17612.Purified recombinant fragment of human PRKACA (AA: 1-120) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	40.6kDa

研究背景 (Background)

cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-

产品名称: PRKACA Mouse Monoclonal Antibody
产品货号: AMM81371




dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. The protein encoded by this gene is a member of the Ser/Thr protein kinase family and is a catalytic subunit of cAMP-dependent protein kinase. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

研究领域 (Research Area)

MAPK signaling pathway

图片 (Image Data)



Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);

注意事项 (Note)

For research use only .