

产品名称: TTR Mouse Monoclonal Antibody
产品货号: AMM81521



产品概述 (Summary)

产品名称 (Production Name)	TTR Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	ELISA,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TTR
别名 (Alternative Names)	CTS; CTS1; PALB; TBPA; HEL111; HsT2651
基因 ID (Gene ID)	7276.0
蛋白 ID (SwissProt ID)	P02766.Purified recombinant fragment of human TTR (AA: 1-147) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	16kDa

研究背景 (Background)

This gene encodes transthyretin, one of the three prealbumins including alpha-1-antitrypsin, transthyretin and


产品名称: TTR Mouse Monoclonal Antibody
产品货号: AMM81521




orosomucoid. Transthyretin is a carrier protein; it transports thyroid hormones in the plasma and cerebrospinal fluid, and also transports retinol (vitamin A) in the plasma. The protein consists of a tetramer of identical subunits. More than 80 different mutations in this gene have been reported; most mutations are related to amyloid deposition, affecting predominantly peripheral nerve and/or the heart, and a small portion of the gene mutations is non-amyloidogenic. The diseases caused by mutations include amyloidotic polyneuropathy, euthyroid hyperthyroxinaemia, amyloidotic vitreous opacities, cardiomyopathy, oculoleptomeningeal amyloidosis, meningocerebrovascular amyloidosis, carpal tunnel syndrome, etc.

研究领域 (Research Area)

图片 (Image Data)



Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);



Flow cytometric analysis of HepG2 cells using TTR mouse mAb (green) and negative control (red).

注意事项 (Note)

For research use only .