

产品名称: RAN Mouse Monoclonal Antibody
产品货号: AMM81657



产品概述 (Summary)

产品名称 (Production Name)	RAN Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ELISA,FC
种属反应性 (Reactivity)	Human,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	RAN
别名 (Alternative Names)	TC4; Gsp1; ARA24
基因 ID (Gene ID)	5901.0
蛋白 ID (SwissProt ID)	P62826.Purified recombinant fragment of human RAN (AA: 1-216) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	24.4kDa

研究背景 (Background)

RAN (ras-related nuclear protein) is a small GTP binding protein belonging to the RAS superfamily that is essential for the


产品名称: RAN Mouse Monoclonal Antibody
产品货号: AMM81657




translocation of RNA and proteins through the nuclear pore complex. The RAN protein is also involved in control of DNA synthesis and cell cycle progression. Nuclear localization of RAN requires the presence of regulator of chromosome condensation 1 (RCC1). Mutations in RAN disrupt DNA synthesis. Because of its many functions, it is likely that RAN interacts with several other proteins. RAN regulates formation and organization of the microtubule network independently of its role in the nucleus-cytosol exchange of macromolecules. RAN could be a key signaling molecule regulating microtubule polymerization during mitosis. RCC1 generates a high local concentration of RAN-GTP around chromatin which, in turn, induces the local nucleation of microtubules. RAN is an androgen receptor (AR) coactivator that binds differentially with different lengths of polyglutamine within the androgen receptor. Polyglutamine repeat expansion in the AR is linked to Kennedy's disease (X-linked spinal and bulbar muscular atrophy). RAN coactivation of the AR diminishes with polyglutamine expansion within the AR, and this weak coactivation may lead to partial androgen insensitivity during the development of Kennedy's disease.

研究领域 (Research Area)

图片 (Image Data)




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using RAN mouse mAb against A431 (1), C6 (2), Jurkat (3), Hela (4), COS7 (5), and Jurkat (6) cell lysate.


产品名称: RAN Mouse Monoclonal Antibody
产品货号: AMM81657



Flow cytometric analysis of Hela cells using RAN mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human esophageal cancer tissues using RAN mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human rectum cancer tissues using RAN mouse mAb with DAB staining.

注意事项 (Note)

For research use only .