产品名称: CD274 Mouse Monoclonal Antibody

产品货号: AMM81688



产品概述 (Summary)

产品名称 (Production Name) CD274 Mouse Monoclonal Antibody

描述 (Description) Mouse monoclonal Antibody

宿主 (Host)Mouse应用 (Application)ELISA,FC种属反应性 (Reactivity)Human

产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG2b 充隆 (Clonality) Monoclonal Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Purified antibody in PBS with 0.05% sodium azide

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CD274

别名 (Alternative Names) B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1

基因 ID (Gene ID) 29126.0

Q9NZQ7.Purified recombinant fragment of human CD274 (AA: 24-153) 蛋白ID (SwissProt ID)

expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio) ELISA 1:5000-1:20000,FC 1:200-1:400

蛋白分子量 (Molecular Weight) 33.3kDa

研究背景 (Background)

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells,

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CD274 Mouse Monoclonal Antibody

产品货号: AMM81688




such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.


The encoded protein is a type I transmembrane protein that has a type I transm

研究领域 (Research Area)

图片 (Image Data)



Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)



Flow cytometric analysis of Hela cells using CD274 mouse mAb (green) and negative control (red).

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838