

产品名称: PSMC3 Mouse Monoclonal Antibody
产品货号: AMM81702



产品概述 (Summary)

产品名称 (Production Name)	PSMC3 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ICC,ELISA,FC
种属反应性 (Reactivity)	Human,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PSMC3
别名 (Alternative Names)	TBP1
基因 ID (Gene ID)	5702.0
蛋白 ID (SwissProt ID)	P17980 .Purified recombinant fragment of human *** (AA: 53-152) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:200-1:1000,ICC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	49kDa

研究背景 (Background)


产品名称: PSMC3 Mouse Monoclonal Antibody
产品货号: AMM81702




The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. A pseudogene has been identified on chromosome 9.

研究领域 (Research Area)

图片 (Image Data)




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using PSMC3 mouse mAb against MCF-7 (1), PC-3 (2), T47D (3), SW620 (4), COS7 (5), C6 (6), HELA (7), and A431 (8) cell lysate.


产品名称: PSMC3 Mouse Monoclonal Antibody
产品货号: AMM81702



Immunofluorescence analysis of MCF-7 cells using PSMC3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)




Immunofluorescence analysis of SK-OV-3 cells using PSMC3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)




Flow cytometric analysis of Hela cells using PSMC3 mouse mAb (green) and negative control (red).

产品名称: PSMC3 Mouse Monoclonal Antibody
产品货号: AMM81702



Immunohistochemical analysis of paraffin-embedded human bladder cancer tissues using PSMC3 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissues using PSMC3 mouse mAb with DAB staining.

注意事项 (Note)

For research use only .