产品名称: NR1I2 Mouse Monoclonal Antibody

产品货号: AMM81774



# 产品概述 (Summary)

产品名称 (Production Name) NR1I2 Mouse Monoclonal Antibody

描述 (**Description**) Mouse monoclonal Antibody

宿主 (Host)Mouse应用 (Application)ELISA,FC种属反应性 (Reactivity)Human

## 产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG1 充隆 (Clonality) Monoclonal Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Purified antibody in PBS with 0.05% sodium azide

纯化方式 (Purification) Affinity Purification

#### 免疫原信息 (Immunogen)

基因名 (Gene Name) NR112

别名 (Alternative Names) BXR; PAR; PRR; PXR; SAR; SXR; ONR1; PAR1; PAR2; PARq

基因 ID (Gene ID) 8856.0

O75469.Purified recombinant fragment of human NR1I2 (AA:1-142) expressed 蛋白 ID (SwissProt ID)

in E. Coli.

### 产品应用 (Application)

稀释比 (Dilution Ratio) ELISA 1:5000-1:20000,FC 1:200-1:400

蛋白分子量 (Molecular Weight) 49.8kDa

# 研究背景 (Background)

This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: NR1I2 Mouse Monoclonal Antibody

产品货号: AMM81774




a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized.<br/>


In the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin.

#### 研究领域 (Research Area)

# 图片 (Image Data)



Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)



Flow cytometric analysis of HepG2 cells using NR112 mouse mAb (green) and negative control (red).

#### 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838