

产品名称: CD50 Mouse Monoclonal Antibody
产品货号: AMM82054



产品概述 (Summary)

产品名称 (Production Name)	CD50 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	ELISA,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CD50
别名 (Alternative Names)	ICAM3; CDW50; ICAM-R
基因 ID (Gene ID)	3385.0
蛋白 ID (SwissProt ID)	P32942.Purified recombinant fragment of human CD50 (AA: extra 30-203) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	59.5kDa

研究背景 (Background)

The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are


产品名称: CD50 Mouse Monoclonal Antibody
产品货号: AMM82054




type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also as a potent signalling molecule. Alternative splicing results in multiple transcript variants encoding different isoforms.

研究领域 (Research Area)

图片 (Image Data)



Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)



Flow cytometric analysis of HL-60 cells using CD50 mouse mAb (green) and negative control (red).

注意事项 (Note)

For research use only .