产品名称: CD158D Mouse Monoclonal Antibody

产品货号: AMM82186



## 产品概述 (Summary)

产品名称 (Production Name) CD158D Mouse Monoclonal Antibody

描述 (**Description**) Mouse monoclonal Antibody

宿主 (Host)Mouse应用 (Application)ELISA,FC种属反应性 (Reactivity)Human

## 产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) Mouse IgG1 充隆 (Clonality) Monoclonal Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Purified antibody in PBS with 0.05% sodium azide

**纯化方式 (Purification)** Affinity Purification

### 免疫原信息 (Immunogen)

基因名 (Gene Name) CD158D

别名 (Alternative Names) KIR2DL4; G9P; KIR103; KIR-2DL4; KIR103AS; KIR-103AS

基因 ID (Gene ID) 3805.0

Q99706.Purified recombinant fragment of human CD158D (AA: extra 22-120) 蛋白 ID (SwissProt ID)

expressed in E. Coli.

## 产品应用 (Application)

稀释比 (Dilution Ratio) ELISA 1:5000-1:20000,FC 1:200-1:400

蛋白分子量 (Molecular Weight) 41.5kDa

### 研究背景 (Background)

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838




产品货号: AMM82186




subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. This gene is one of the "framework" loci that is present on all haplotypes. Alternate alleles of this gene are represented on multiple alternate reference loci (ALT\_REF\_LOCs). Alternative splicing results in multiple transcript variants, some of which may not be annotated on the primary reference assembly.

#### 研究领域 (Research Area)

## 图片 (Image Data)



Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)



Flow cytometric analysis of HL-60 cells using CD158D mouse mAb (green) and negative control (red).

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: CD158D Mouse Monoclonal Antibody

产品货号: AMM82186



# 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838