

产品名称: ATXN1 Mouse Monoclonal Antibody
产品货号: AMM82210



产品概述 (Summary)

产品名称 (Production Name)	ATXN1 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ELISA,FC
种属反应性 (Reactivity)	Human,Mouse,Rat, Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ATXN1
别名 (Alternative Names)	ATX1; SCA1; D6S504E
基因 ID (Gene ID)	6310.0
蛋白 ID (SwissProt ID)	P54253.Purified recombinant fragment of human ATXN1 (AA: 645-815) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:500,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	86.9kDa

研究背景 (Background)

The autosomal dominant cerebellar ataxias (ADCA) are a heterogeneous group of neurodegenerative disorders


产品名称: ATXN1 Mouse Monoclonal Antibody
产品货号: AMM82210




characterized by progressive degeneration of the cerebellum, brain stem and spinal cord. Clinically, ADCA has been divided into three groups: ADCA types I-III. ADCAI is genetically heterogeneous, with five genetic loci, designated spinocerebellar ataxia (SCA) 1, 2, 3, 4 and 6, being assigned to five different chromosomes. ADCAII, which always presents with retinal degeneration (SCA7), and ADCAIII often referred to as the 'pure' cerebellar syndrome (SCA5), are most likely homogeneous disorders. Several SCA genes have been cloned and shown to contain CAG repeats in their coding regions. ADCA is caused by the expansion of the CAG repeats, producing an elongated polyglutamine tract in the corresponding protein. The expanded repeats are variable in size and unstable, usually increasing in size when transmitted to successive generations. The function of the ataxins is not known. This locus has been mapped to chromosome 6, and it has been determined that the diseased allele contains 40-83 CAG repeats, compared to 6-39 in the normal allele, and is associated with spinocerebellar ataxia type 1 (SCA1). Alternative splicing results in multiple transcript variants, with one variant encoding multiple distinct proteins, ATXN1 and Alt-ATXN1, due to the use of overlapping alternate reading frames.

研究领域 (Research Area)

图片 (Image Data)




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using ATXN1 mouse mAb against C6 (1), COS7 (2), NIH/3T3 (3), and HL-60 (4) cell lysate.


产品名称: ATXN1 Mouse Monoclonal Antibody
产品货号: AMM82210



Western blot analysis using ATXN1 mouse mAb against F9(1)L1210(2)C2C12(3) cell lysate.




Flow cytometric analysis of Jurkat cells using ATXN1 mouse mAb (green) and negative control (red).




Immunohistochemical analysis of paraffin-embedded human endometrial cancer tissues using ATXN1 mouse mAb with DAB staining.

产品名称: ATXN1 Mouse Monoclonal Antibody
产品货号: AMM82210



Immunohistochemical analysis of paraffin-embedded Mouse brain(A) Mouse kidney(B) using ATXN1 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded Rat brain(A) Rat kidney(B) using ATXN1 mouse mAb with DAB staining.

注意事项 (Note)

For research use only .