

产品名称: CCNE1 Mouse Monoclonal Antibody
产品货号: AMM82445



产品概述 (Summary)

产品名称 (Production Name)	CCNE1 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	IHC,ELISA,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CCNE1
别名 (Alternative Names)	CCNE; Pccne1
基因 ID (Gene ID)	898.0
蛋白 ID (SwissProt ID)	P24864.Purified recombinant fragment of human CCNE1 (AA: 1-100) expressed in E. Coli.

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
蛋白分子量 (Molecular Weight)	47kDa

研究背景 (Background)

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a

产品名称: CCNE1 Mouse Monoclonal Antibody
产品货号: AMM82445




dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. [provided by RefSeq, Apr 2016]


研究领域 (Research Area)

PI3K-Akt signaling pathway, mTOR signaling pathway

图片 (Image Data)




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)



Flow cytometric analysis of HeLa cells using CCNE1 mouse mAb (green) and negative control (red).

产品名称: CCNE1 Mouse Monoclonal Antibody
产品货号: AMM82445



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissues using CCNE1 mouse mAb with DAB staining.

注意事项 (Note)

For research use only .