

产品名称: CD5 Mouse Monoclonal Antibody
产品货号: AMM84944



产品概述 (Summary)

产品名称 (Production Name)	CD5 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	IHC
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide,0.5%protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CD5
别名 (Alternative Names)	CD5; LEU1; T-cell surface glycoprotein CD5; Lymphocyte antigen T1/Leu-1; CD antigen CD5
基因 ID (Gene ID)	921.0
蛋白 ID (SwissProt ID)	P06127.Synthetic peptide conjugated to KLH.

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:100
蛋白分子量 (Molecular Weight)	/

研究背景 (Background)


产品名称: CD5 Mouse Monoclonal Antibody
产品货号: AMM84944




May act as a receptor in regulating T-cell proliferation. CD5 interacts with CD72/LYB-2.

研究领域 (Research Area)


图片 (Image Data)



Immunofluorescence analysis of CD5 in mouse heart tissue using CD5 antibody(10G8)(red),and DAPI (blue).




Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Human tonsils using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.

产品名称: CD5 Mouse Monoclonal Antibody
产品货号: AMM84944



Immunohistochemistry analysis of paraffin-embedded mouse kidney tissue using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunohistochemistry analysis of paraffin-embedded Human Tonsil Carcinoma using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

注意事项 (Note)

For research use only .