

产品名称: FGFR1 Mouse Monoclonal Antibody
产品货号: AMM85972



产品概述 (Summary)

产品名称 (Production Name)	FGFR1 Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,IHC,ICC
种属反应性 (Reactivity)	Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FGFR1
别名 (Alternative Names)	Fibroblast growth factor receptor 1, FGFR-1, Basic fibroblast growth factor receptor 1, BFGFR, bFGF-R-1, Fms-like tyrosine kinase 2, FLT-2, N-sam, Proto-oncogene c-Fgr, CD331, FGFR1, BFGFR, CEK, FGFR, FLG, FLT2, HBGFR
基因 ID (Gene ID)	2260.0
蛋白 ID (SwissProt ID)	P11362. This FGFR1 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 806-842 amino acids from the C-terminal region of human FGFR1.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, IHC 1:100-1:500, ICC 1:25-1:50
蛋白分子量 (Molecular Weight)	91.9kDa

产品名称: FGFR1 Mouse Monoclonal Antibody
产品货号: AMM85972




研究背景 (Background)

Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of embryonic development, cell proliferation, differentiation and migration. Required for normal mesoderm patterning and correct axial organization during embryonic development, normal skeletogenesis and normal development of the gonadotropin-releasing hormone (GnRH) neuronal system. Phosphorylates PLCG1, FRS2, GAB1 and SHB. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes phosphorylation of SHC1, STAT1 and PTPN11/SHP2. In the nucleus, enhances RPS6KA1 and CREB1 activity and contributes to the regulation of transcription. FGFR1 signaling is down-regulated by IL17RD/SEF, and by FGFR1 ubiquitination, internalization and degradation.


研究领域 (Research Area)

TGF-beta signaling pathway, PI3K-Akt signaling pathway, MAPK signaling pathway, Hippo signaling pathway

图片 (Image Data)




Western blot analysis of lysate from HeLa cell line, using FGFR1 Antibody (C-term)(Cat. #AMM85972). AMM85972 was diluted at 1:2000. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 20µg.



产品名称: FGFR1 Mouse Monoclonal Antibody
产品货号: AMM85972



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling FGFR1 with AMM85972 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on HeLa cell line. The nuclear counter stain is DAPI (blue).



AMM85972 staining FGFR1 in human brain tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

注意事项 (Note)

For research use only .