

产品名称: TERF2IP Mouse Monoclonal Antibody
产品货号: AMM86038



产品概述 (Summary)

产品名称 (Production Name)	TERF2IP Mouse Monoclonal Antibody
描述 (Description)	Mouse monoclonal Antibody
宿主 (Host)	Mouse
应用 (Application)	WB,FC
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	Mouse IgG1
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TERF2IP
别名 (Alternative Names)	Telomeric repeat-binding factor 2-interacting protein 1, TERF2-interacting telomeric protein 1, TRF2-interacting telomeric protein 1, Dopamine receptor-interacting protein 5, Repressor/activator protein 1 homolog, RAP1 homolog, hRap1, TERF2IP, DRIP5, RAP1
基因 ID (Gene ID)	54386.0
蛋白 ID (SwissProt ID)	Q9NYB0. This TERF2IP antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between amino acids from the human region of human TERF2IP.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000, FC 1:25-1:50
----------------------	--------------------------------

产品名称: TERF2IP Mouse Monoclonal Antibody
产品货号: AMM86038




蛋白分子量 (Molecular Weight) 44.2kDa

研究背景 (Background)

Acts both as a regulator of telomere function and as a transcription regulator. Involved in the regulation of telomere length and protection as a component of the shelterin complex (telosome). In contrast to other components of the shelterin complex, it is dispensable for telomere capping and does not participate in the protection of telomeres against non-homologous end-joining (NHEJ)-mediated repair. Instead, it is required to negatively regulate telomere recombination and is essential for repressing homology-directed repair (HDR), which can affect telomere length. Does not bind DNA directly: recruited to telomeric double-stranded 5'-TTAGGG-3' repeats via its interaction with TERF2. Independently of its function in telomeres, also acts as a transcription regulator: recruited to extratelomeric 5'-TTAGGG-3' sites via its association with TERF2 or other factors, and regulates gene expression. When cytoplasmic, associates with the I-kappa-B-kinase (IKK) complex and acts as a regulator of the NF-kappa-B signaling by promoting IKK-mediated phosphorylation of RELA/p65, leading to activate expression of NF-kappa-B target genes.

研究领域 (Research Area)

图片 (Image Data)



All lanes : Anti-TERF2IP Antibody at 1:2000 dilution

注意事项 (Note)

For research use only .