

产品名称: Cyclin E2 Rabbit Polyclonal Antibody
产品货号: APRab00065



产品概述 (Summary)

产品名称 (Production Name)	Cyclin E2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
纯化方式 (Purification)	Affinity Chromatography

免疫原信息 (Immunogen)

基因名 (Gene Name)	CCNE2
别名 (Alternative Names)	CCNE2; G1/S-specific cyclin-E2
基因 ID (Gene ID)	9134
蛋白 ID (SwissProt ID)	O96020.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000,IHC 1:50-1:100,ICC/IF 1:50-1:200,IP 1:20-1:50
蛋白分子量 (Molecular Weight)	Calculated MW: 47 kDa; Observed MW: 50 kDa

研究背景 (Background)

The human Cyclin E2 gene encodes a 404 amino acid protein that is most closely related to Cyclin E. Cyclin E2 mRNA levels

产品名称: Cyclin E2 Rabbit Polyclonal Antibody
产品货号: AP Rab00065




peaks at the G1 / S transition. Cyclin E2 associates with Cdk2 in a functional kinase complex that is inhibited by both p27 (Kip1) and p21 (Cip1). Cyclin E2 / Cdk2 phosphorylates histone H1 in vitro. G1 cyclin E controls the initiation of DNA synthesis by activating CDK2. Abnormally high levels of cyclin E expression have frequently been observed in human cancers.

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western blot analysis of Cyclin E2 in Jurkat lysates using Cyclin E2 antibody.

注意事项 (Note)

For research use only .