

产品名称: 5-LO (phospho Ser272) Rabbit Polyclonal Antibody
产品货号: APRab04192



产品概述 (Summary)

产品名称 (Production Name)	5-LO (phospho Ser272) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ALOX5
别名 (Alternative Names)	ALOX5; LOG5; Arachidonate 5-lipoxygenase; 5-LO; 5-lipoxygenase
基因 ID (Gene ID)	240.0
蛋白 ID (SwissProt ID)	P09917.The antiserum was produced against synthesized peptide derived from human Arachidonate 5 Lipoxygenase around the phosphorylation site of Ser271. AA range:246-295

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	78kDa

产品名称: 5-LO (phospho Ser272) Rabbit Polyclonal Antibody
产品货号: APRab04192




研究背景 (Background)

This gene encodes a member of the lipoxygenase gene family and plays a dual role in the synthesis of leukotrienes from arachidonic acid. The encoded protein, which is expressed specifically in bone marrow-derived cells, catalyzes the conversion of arachidonic acid to 5(S)-hydroperoxy-6-trans-8,11,14-cis-eicosatetraenoic acid, and further to the allylic epoxide 5(S)-trans-7,9-trans-11,14-cis-eicosatetraenoic acid (leukotriene A4). Leukotrienes are important mediators of a number of inflammatory and allergic conditions. Mutations in the promoter region of this gene lead to a diminished response to antileukotriene drugs used in the treatment of asthma and may also be associated with atherosclerosis and several cancers. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012], catalytic activity: Arachidonate + O(2) = leukotriene A(4) + H(2)O., cofactor: Binds 1 iron ion per subunit., cofactor: Binds 2 calcium ions per subunit., function: Catalyzes the first step in leukotriene biosynthesis, and thereby plays a role in inflammatory processes., pathway: Lipid metabolism; leukotriene A4 biosynthesis., PTM: Serine phosphorylation by MAPKAPK2 is stimulated by arachidonic acid. Phosphorylation on Ser-523 by PKA has an inhibitory effect. Phosphorylation on Ser-272 prevents export from the nucleus., similarity: Belongs to the lipoxygenase family., similarity: Contains 1 lipoxygenase domain., similarity: Contains 1 PLAT domain., subcellular location: Shuttles between cytoplasm and nucleus. Found exclusively in the nucleus, when phosphorylated on Ser-272. Calcium binding promotes translocation from the cytosol and the nuclear matrix to the nuclear envelope and membrane association., subunit: Interacts with ALOX5AP and LTC4S.,

研究领域 (Research Area)


Arachidonic acid metabolism;

图片 (Image Data)




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Arachidonate 5 Lipoxygenase (Phospho-Ser271) Antibody

产品名称: 5-LO (phospho Ser272) Rabbit Polyclonal Antibody
产品货号: APRab04192



Immunohistochemistry analysis of paraffin-embedded human skeletal muscle, using Arachidonate 5 Lipxygenase (Phospho-Ser271) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HUVEC cells, using Arachidonate 5 Lipxygenase (Phospho-Ser271) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .