

产品名称: APLP2 (phospho Tyr755) Rabbit Polyclonal Antibody
产品货号: APRab04247



产品概述 (Summary)

产品名称 (Production Name)	APLP2 (phospho Tyr755) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	APLP2
别名 (Alternative Names)	APLP2; APPL2; Amyloid-like protein 2; APLP-2; APPH; Amyloid protein homolog; CDEI box-binding protein; CDEBP
基因 ID (Gene ID)	334.0
蛋白 ID (SwissProt ID)	Q06481. The antiserum was produced against synthesized peptide derived from human APLP2 around the phosphorylation site of Tyr755. AA range: 714-763

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:10000-1:20000
----------------------	---

产品名称: APLP2 (phospho Tyr755) Rabbit Polyclonal Antibody
产品货号: APRab04247



蛋白分子量 (Molecular Weight)


研究背景 (Background)

This gene encodes amyloid precursor- like protein 2 (APLP2), which is a member of the APP (amyloid precursor protein) family including APP, APLP1 and APLP2. This protein is ubiquitously expressed. It contains heparin-, copper- and zinc-binding domains at the N-terminus, BPTI/Kunitz inhibitor and E2 domains in the middle region, and transmembrane and intracellular domains at the C-terminus. This protein interacts with major histocompatibility complex (MHC) class I molecules. The synergy of this protein and the APP is required to mediate neuromuscular transmission, spatial learning and synaptic plasticity. This protein has been implicated in the pathogenesis of Alzheimer's disease. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Aug 2011], alternative products: Additional isoforms seem to exist, function: May play a role in the regulation of hemostasis. The soluble form may have inhibitory properties towards coagulation factors. May interact with cellular G-protein signaling pathways. May bind to the DNA 5'-GTCACATG-3' (CDEI box). Inhibits trypsin, chymotrypsin, plasmin, factor XIA and plasma and glandular kallikrein., PTM: The BPTI/Kunitz inhibitor domain is O-glycosylated., similarity: Belongs to the APP family., similarity: Contains 1 BPTI/Kunitz inhibitor domain., subunit: Interacts with CPEB1., tissue specificity: In placenta, brain, heart, lung, liver, kidney and endothelial tissues.,

研究领域 (Research Area)

Neuroscience

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain, using APLP2 (Phospho-Tyr755) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .