

产品货号: APRab04262

产品概述 (Summary)

产品名称 (Production Name) Arrestin-β-1 (phospho Ser412) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Monkey

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) ARRB1

别名 (Alternative Names) ARRB1; ARR1; Beta-arrestin-1; Arrestin beta-1

基因 ID (Gene ID) 408.0

P49407.The antiserum was produced against synthesized peptide derived

蛋白ID (SwissProt ID) from human Arrestin 1 around the phosphorylation site of Ser412. AA

range:369-418

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 47kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838





产品货号: APRab04262




研究背景 (Background)

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of Gprotein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011], function: Regulates beta-adrenergic receptor function. Beta-arrestins seem to bind phosphorylated beta-adrenergic receptors, thereby causing a significant impairment of their capacity to activate G(S) proteins, online information: Arrestin entry, similarity: Belongs to the arrestin family,


研究领域(Research Area)

MAPK ERK Growth; MAPK G Protein; Chemokine; Endocytosis;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Arrestin 1 (Phospho-Ser412) Antibody. The picture on the right is blocked with the phospho peptide.




E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



产品货号: APRab04262

Western blot analysis of lysates from COS7 cells treated with Etoposide 25uM 60 ', using Arrestin 1 (Phospho-Ser412)

Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838