

产品名称: AR-β2 (phospho Ser355/S356) Rabbit Polyclonal Antibody
产品货号: APRab04265



产品概述 (Summary)

产品名称 (Production Name)	AR-β2 (phospho Ser355/S356) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ADRB2
别名 (Alternative Names)	ADRB2; ADRB2R; B2AR; Beta-2 adrenergic receptor; Beta-2 adrenoreceptor; Beta-2 adrenoceptor
基因 ID (Gene ID)	154.0
蛋白 ID (SwissProt ID)	P07550.The antiserum was produced against synthesized peptide derived from human Adrenergic Receptor B2 around the phosphorylation site of Ser355 and Ser356. AA range:331-380

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
----------------------	--

产品名称: AR- β 2 (phospho Ser355/S356) Rabbit Polyclonal Antibody
产品货号: APRab04265

蛋白分子量 (Molecular Weight) 47kDa


研究背景 (Background)

This gene encodes beta-2-adrenergic receptor which is a member of the G protein-coupled receptor superfamily. This receptor is directly associated with one of its ultimate effectors, the class C L-type calcium channel Ca(V)1.2. This receptor-channel complex also contains a G protein, an adenylyl cyclase, cAMP-dependent kinase, and the counterbalancing phosphatase, PP2A. The assembly of the signaling complex provides a mechanism that ensures specific and rapid signaling by this G protein-coupled receptor. This gene is intronless. Different polymorphic forms, point mutations, and/or downregulation of this gene are associated with nocturnal asthma, obesity and type 2 diabetes. [provided by RefSeq, Jul 2008],disease:Polymorphic forms of ADRB2 could impart some form of nocturnal asthma.,function:Beta-adrenergic receptors mediate the catecholamine-induced activation of adenylate cyclase through the action of G proteins. The beta-2-adrenergic receptor binds epinephrine with an approximately 30-fold greater affinity than it does norepinephrine.,PTM:Palmitoylated; may reduce accessibility of Ser-345 and Ser-346 by anchoring Cys-341 to the plasma membrane. Agonist stimulation promotes depalmitoylation and further allows Ser-345 and Ser-346 phosphorylation.,PTM:Phosphorylated by PKA and BARK upon agonist stimulation, which mediates homologous desensitization of the receptor. PKA-mediated phosphorylation seems to facilitate phosphorylation by BARK. Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Phosphorylation of Tyr-141 is induced by insulin and leads to supersensitization of the receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,subunit:Bounds SLC9A3R1 and GPRASP1.,

研究领域 (Research Area)


Calcium;Neuroactive ligand-receptor interaction;Endocytosis;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain, using Adrenergic Receptor B2 (Phospho-Ser355+Ser356) Antibody. The picture on the right is blocked with the phospho peptide.

产品名称: AR- β 2 (phospho Ser355/S356) Rabbit Polyclonal Antibody
产品货号: APRab04265



Western blot analysis of lysates from HUVEC cells treated with serum 20% 15 ' and COS7 cells treated with serum 20% 15 ', using Adrenergic Receptor B2 (Phospho-Ser355+Ser356) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .