

产品名称: ATF-2 (phospho Ser472) Rabbit Polyclonal Antibody
产品货号: APRab04274



产品概述 (Summary)

产品名称 (Production Name)	ATF-2 (phospho Ser472) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ATF2
别名 (Alternative Names)	ATF2; CREB2; CREBP1; Cyclic AMP-dependent transcription factor ATF-2; cAMP-dependent transcription factor ATF-2; Activating transcription factor 2; Cyclic AMP-responsive element-binding protein 2; CREB-2; cAMP-responsive element-binding pro
基因 ID (Gene ID)	1386.0
蛋白 ID (SwissProt ID)	P15336. The antiserum was produced against synthesized peptide derived from human ATF-2 around the phosphorylation site of Ser472. AA range: 441-490

产品应用 (Application)

产品名称: ATF-2 (phospho Ser472) Rabbit Polyclonal Antibody
产品货号: APRab04274



稀释比 (Dilution Ratio)

IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight)


研究背景 (Background)

activating transcription factor 2(ATF2) Homo sapiens This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription. This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. The encoded protein may also be involved in cell's DNA damage response independent of its role in transcriptional regulation. Several alternatively spliced transcript variants have been found for this gene [provided by RefSeq, Jan 2014] caution: It is uncertain whether Met-1 or Met-19 is the initiator., function: Transcriptional activator, probably constitutive, which binds to the cAMP-responsive element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Interaction with JUN redirects JUN to bind to CREs preferentially over the 12-O-tetradecanoylphorbol-13-acetate response elements (TREs) as part of an ATF2-c-Jun complex., PTM: Phosphorylation of Thr-69 and Thr-71 by MAPK14 causes increased transcriptional activity. Also phosphorylated and activated by JNK., similarity: Belongs to the bZIP family., similarity: Belongs to the bZIP family. ATF subfamily., similarity: Contains 1 bZIP domain., similarity: Contains 1 C2H2-type zinc finger., subunit: Binds DNA as a dimer and can form a homodimer in the absence of DNA. Can form a heterodimer with JUN. Interacts with SMAD3 and SMAD4. Binds through its N-terminal region to UTF1 which acts as a coactivator of ATF2 transcriptional activity., tissue specificity: Abundant expression seen in the brain.,

研究领域 (Research Area)

B Cell Receptor; Stem cell pathway; MAPK_ERK_Growth; MAPK_G_Protein; Akt_PKB; Protein_Acetylation


图片 (Image Data)



产品名称: ATF-2 (phospho Ser472) Rabbit Polyclonal Antibody
产品货号: APRab04274



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using ATF-2 (Phospho-Ser472) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using ATF-2 (Phospho-Ser472) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .