

产品名称: Axl (phospho-Tyr702) Rabbit Polyclonal Antibody
产品货号: APRab04290



产品概述 (Summary)

产品名称 (Production Name)	Axl (phospho-Tyr702) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	AXL
别名 (Alternative Names)	AXL; UFO; Tyrosine-protein kinase receptor UFO; AXL oncogene
基因 ID (Gene ID)	558.0
蛋白 ID (SwissProt ID)	P30530.Synthesized phosho peptide around human Axl (Tyr702)

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000
蛋白分子量 (Molecular Weight)	130kDa

研究背景 (Background)

The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation


产品名称: Axl (phospho-Tyr702) Rabbit Polyclonal Antibody
产品货号: AP Rab04290

and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a protein = ADP + a phosphoprotein., caution: Although authors have considered STK6 and STK15 as two different proteins, it is clear that they are the same protein., disease: Defects in AURKA are responsible for numerical centrosome aberrations including aneuploidy., function: May play a role in cell cycle regulation during anaphase and/or telophase, in relation to the function of the centrosome/spindle pole region during chromosome segregation. May be involved in microtubule formation and/or stabilization. Phosphorylates ARHGEF2 and BORA., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. Aurora subfamily., similarity: Contains 1 protein kinase domain., subcellular location: Localizes on centrosomes in interphase cells and at each spindle pole in mitosis., subunit: Interacts with TACC1 and CPEB1. Interacts with its substrates BORA and ARHGEF2., tissue specificity: Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines. Expression is cell-cycle regulated, low in G1/S, accumulates during G2/M, and decreases rapidly after.,

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western Blot analysis of HeLa treated or untreated by LPS lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only .