

产品名称: Brk (phospho Tyr447) Rabbit Polyclonal Antibody
产品货号: AP Rab04334



产品概述 (Summary)

产品名称 (Production Name)	Brk (phospho Tyr447) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PTK6
别名 (Alternative Names)	PTK6; BRK; Protein-tyrosine kinase 6; Breast tumor kinase; Tyrosine-protein kinase BRK
基因 ID (Gene ID)	5753.0
蛋白 ID (SwissProt ID)	Q13882. The antiserum was produced against synthesized peptide derived from human Breast Tumor Kinase around the phosphorylation site of Tyr447. AA range: 402-451

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, ICC/IF 1:200-1:1000, ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	50kDa

产品名称: Brk (phospho Tyr447) Rabbit Polyclonal Antibody
产品货号: APRab04334




研究背景 (Background)

protein tyrosine kinase 6 (PTK6) Homo sapiens The protein encoded by this gene is a cytoplasmic nonreceptor protein kinase which may function as an intracellular signal transducer in epithelial tissues. Overexpression of this gene in mammary epithelial cells leads to sensitization of the cells to epidermal growth factor and results in a partially transformed phenotype. Expression of this gene has been detected at low levels in some breast tumors but not in normal breast tissue. The encoded protein has been shown to undergo autophosphorylation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2012], catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate, enzyme regulation: Activated enzyme seems to have greater access to its substrates, function: Phosphorylates KHDRBS1, KHDRBS2, KHDRBS3 and STAP2/BKS. May function as an intracellular signal transducer in epithelial tissues. Overexpression in mammary cells leads to mitogenically sensitization to EGF, and results in a partially transformed phenotype. Its presence in the nucleus appears to be linked to suppression of tumor progression, PTM: Autophosphorylated. The phosphorylation of Tyr-447 may lead to the autoinhibition of the enzyme, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. BRK/PTK6/SIK subfamily, similarity: Contains 1 protein kinase domain, similarity: Contains 1 SH2 domain, similarity: Contains 1 SH3 domain, subcellular location: Colocalizes with KHDRBS1, KHDRBS2 or KHDRBS3, within the nucleus. In secretory epithelial cells from prostate adenocarcinoma, nuclear localization is higher in low-grade and lower in high-grade regions of the tumors, subunit: Interacts with GAP-A.p65 (By similarity). Interacts with KHDRBS1. Interacts with phosphorylated IRS4, tissue specificity: Epithelia-specific. Very high level in colon and high levels in small intestine and prostate, and low levels in some fetal tissues. Expressed at low level in some breast tumors, but not in normal breast. Also found in melanocytes. Not expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas,

研究领域 (Research Area)

图片 (Image Data)




Immunofluorescence analysis of NIH/3T3 cells, using Breast Tumor Kinase (Phospho-Tyr447) Antibody. The picture on the


产品名称: Brk (phospho Tyr447) Rabbit Polyclonal Antibody
产品货号: AP Rab04334



right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with EGF 200ng/ml 30', using Breast Tumor Kinase (Phospho-Tyr447) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of 293 cells using Phospho-Brk (Y447) Polyclonal Antibody

注意事项 (Note)

For research use only .