

产品概述 (Summary)

产品名称 (Production Name) C3G (phospho Tyr504) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) RAPGEF1

RAPGEF1; GRF2; Rap guanine nucleotide exchange factor 1; CRK SH3-binding 别名 (Alternative Names)

GNRP; Guanine nucleotide-releasing factor 2; Protein C3G

基因 ID (Gene ID) 2889.0

Q13905.The antiserum was produced against synthesized peptide derived

蛋白ID (SwissProt ID) from human RapGEF1 around the phosphorylation site of Tyr504. AA


range:470-519

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 120kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab04343



研究背景 (Background)


This gene encodes a human guanine nucleotide exchange factor. It transduces signals from CRK by binding the SH3 domain of CRK, and activating several members of the Ras family of GTPases. This signaling cascade that may be involved in apoptosis, integrin-mediated signal transduction, and cell transformation. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some variants has not been determined. [provided by RefSeq, Jul 2008], function: Guanine nucleotide-releasing protein that binds to SH3 domain of CRK and GRB2/ASH.

Transduces signals from CRK to activate RAS., similarity: Contains 1 N-terminal Ras-GEF domain., similarity: Contains 1 Ras-GEF domain., subunit: Interacts with CRK via its SH3-binding sites., tissue specificity: Ubiquitously expressed in adult and fetus. Expression is high in adult skeletal muscle and placenta and in fetal brain and heart. Low levels of expression in adult and fetal liver.,


研究领域 (Research Area)

Regulation of Actin Dynamics; AMPK

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using RapGEF1 (Phospho-Tyr504) Antibody



Western blot analysis of lysates from HepG2 cells treated with Na3VO4 0.3nM 40 ', using RapGEF1 (Phospho-Tyr504)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838



Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838