

## 产品概述 (Summary)

产品名称 (Production Name) CaMKIIα/δ (phospho Thr286) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse,Rat,Other

# 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

# 免疫原信息 (Immunogen)

别名 (Alternative Names)

基因名 (Gene Name) CAMK2A/CAMK2D

CAMK2A; CAMKA; KIAA0968; Calcium/calmodulin-dependent protein kinase

type II subunit alpha; CaM kinase II subunit alpha; CaMK-II subunit alpha;

CAMK2D; CAMKD; Calcium/calmodulin-dependent protein kinase type II

subunit delta; CaM kinase II

基因 ID (Gene ID) 815/817

Q9UQM7/Q13557.The antiserum was produced against synthesized peptide

蛋白ID (SwissProt ID) derived from human CaMK2 around the phosphorylation site of Thr286. AA

range:256-305

# 产品应用 (Application)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838







WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000

EnkiLife

蛋白分子量 (Molecular Weight) 54kDa

### 研究背景 (Background)


稀释比 (Dilution Ratio)

The product of this gene belongs to the serine/threonine protein kinases family, and to the Ca(2+)/calmodulin-dependent protein kinases subfamily. Calcium signaling is crucial for several aspects of plasticity at glutamatergic synapses. This calcium calmodulin-dependent protein kinase is composed of four different chains: alpha, beta, gamma, and delta. The alpha chain encoded by this gene is required for hippocampal long-term potentiation (LTP) and spatial learning. In addition to its calcium-calmodulin (CaM)-dependent activity, this protein can undergo autophosphorylation, resulting in CaMindependent activity. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Nov 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Autophosphorylation of Thr-286 allows the kinase to switch from a calmodulin-dependent to a calmodulin-independent state., function: CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDARdependent potentiation of the AMPAR and synaptic plasticity, similarity: Belongs to the protein kinase superfamily, similarity: Belongs to the protein kinase superfamily, CAMK Ser/Thr protein kinase family, CaMK subfamily, similarity: Contains 1 protein kinase domain...subcellular location: Postsynaptic lipid rafts...subunit: CAMK2 is composed of four different chains: alpha, beta, gamma, and delta. The different isoforms assemble into homo- or heteromultimeric holoenzymes composed of 8 to 12 subunits. Interacts with BAALC, MPDZ, SYN1, CAMK2N2 and SYNGAP1.,

#### 研究领域 (Research Area)

ErbB\_HER;Calcium;Oocyte meiosis;WNT;WNT-T CELLLong-term potentiation;Neurotrophin;Olfactory transduction;GnRH;Melanogenesis;Glioma;

# 图片 (Image Data)




Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838







Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using CaMK2 (Phospho-Thr286) Antibody



Immunofluorescence analysis of COS7 cells, using CaMK2 (Phospho-Thr286) Antibody. The picture on the right is blocked with the phospho peptide.




Immunohistochemistry analysis of paraffin-embedded human brain, using CaMK2 (Phospho-Thr286) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from K562 cells, using CaMK2 (Phospho-Thr286) Antibody. The lane on the right is blocked with the phospho peptide.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838





Western Blot analysis of HELA cells using Phospho-CaMKII $\alpha/\delta$  (T286) Polyclonal Antibody diluted at 1: 500

# 注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838