

产品货号: APRab04381

产品概述 (Summary)

产品名称 (Production Name) Catenin δ-1 (phospho-Tyr904) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB

种属反应性 (Reactivity) Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CTNND1

Catenin delta-1 (Cadherin-associated Src substrate) (CAS) (p120 catenin) 別名 (Alternative Names)

(p120(ctn)) (p120(cas))

基因 ID (Gene ID) 1500.0

蛋白ID (SwissProt ID) O60716.Synthesized phosho peptide around human Catenin δ-1 (Tyr904)

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:1000-1:2000

蛋白分子量 (Molecular Weight) 108kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

Antibody

产品货号: APRab04381




研究背景 (Background)

catenin delta 1(CTNND1) Homo sapiens This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene. [provided by RefSeq, Dec 2010], alternative products: Experimental confirmation may be lacking for some isoforms, disease: May contribute to cell malignancy. Complete loss of expression was observed in approximately 10% of invasive ductal breast carcinomas investigated.,domain:A possible nuclear localization signal exists in all isoforms where Asp-626--631-Arg are deleted., function: Binds to and inhibits the transcriptional repressor ZBTB33, which may lead to activation of target genes of the Wnt signaling pathway (By similarity). May associate with and regulate the cell adhesion properties of both C- and E-cadherins. Implicated both in cell transformation by SRC and in ligand-induced receptor signaling through the EGF, PDGF, CSF-1 and ERBB2 receptors. Promotes GLIS2 C-terminal cleavage., induction: Induced in vascular endothelium by wounding. This effect is potentiated by prior laminar shear stress, which enhances wound closure., PTM: Phosphorylated., similarity: Belongs to the beta-catenin family., similarity: Contains 10 ARM repeats, subcellular location: Interaction with GLIS2 promotes nuclear translocation, subunit: Belongs to a multiprotein cell-cell adhesion complex that also contains E-cadherin, alpha-catenin, beta-catenin, and gamma-catenin. Binds to the Cterminal fragment of PSEN1 and mutually competes for E-cadherin. Interacts with ZBTB33. Interacts with GLIS2., tissue specificity: Expressed in vascular endothelium.,

研究领域(Research Area)

Adherens Junction; Leukocyte transendothelial migration;

图片 (Image Data)



Western Blot analysis of Hela treated or untreated by LPS lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项(Note)

E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



产品货号: APRab04381

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838