产品名称: CD45 (phospho Ser1007) Rabbit Polyclonal

Antibody

产品货号: APRab04405



产品概述 (Summary)

产品名称 (Production Name) CD45 (phospho Ser1007) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) PTPRC

PTPRC; CD45; Receptor-type tyrosine-protein phosphatase C; Leukocyte 別名 (Alternative Names)

common antigen; L-CA; T200; CD antigen CD45

基因 ID (Gene ID) 5788.0

P08575.The antiserum was produced against synthesized peptide derived

蛋白 ID (SwissProt ID) from human CD45 around the phosphorylation site of Ser1007. AA range:981-

1030

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CD45 (phospho Ser1007) Rabbit Polyclonal

Antibody

产品货号: APRab04405



蛋白分子量 (Molecular Weight) 150kDa

研究背景 (Background)

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encalternative products: At least 8 isoforms are produced, catalytic activity: Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate, disease: Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+)SCID) [MIM:608971]. SCID refers to a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients with SCID present in infancy with recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development, disease: Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain.,domain:The first PTPase domain interacts with SKAP1.,function:Required for T-cell activation through the antigen receptor. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits an dephosphorylates SKAP1 and FYN, online information:CD45 entry, online information: PTPRC mutation db, PTM: Heavily N- and O-glycosylated., similarity: Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily, similarity: Contains 2 fibronectin type-III domains, similarity: Contains 2 tyrosine-protein phosphatase domains., subunit: Binds GANAB and PRKCSH (By similarity). Interacts with SKAP1.,

研究领域 (Research Area)

Cell adhesion molecules (CAMs);T_Cell_Receptor;Fc gamma R-mediated phagocytosis;Primary immunodeficiency;

图片 (Image Data)


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: CD45 (phospho Ser1007) Rabbit Polyclonal


Antibody

产品货号: APRab04405





Immunofluorescence analysis of HeLa cells, using CD45 (Phospho-Ser1007) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with TNF 20ng/ml 15 ', using CD45 (Phospho-Ser1007) Antibody.

The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838