

Antibody

产品货号: APRab04432



产品概述 (Summary)

产品名称 (Production Name) Cdk1/2/3 (phospho Thr14) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Mouse, Rat, Monkey

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) lgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

别名 (Alternative Names)

基因名 (Gene Name) CDK1/CDK2/CDK3

CDK1; CDC2; CDC28A; CDKN1; P34CDC2; Cyclin-dependent kinase 1; CDK1;

Cell division control protein 2 homolog; Cell division protein kinase 1; p34

protein kinase; CDK2; CDKN2; Cyclin-dependent kinase 2; Cell division protein

kinase 2; p33 pr

基因 ID (Gene ID) 1017.0

P06493/P24941/Q00526.The antiserum was produced against synthesized

蛋白 ID (SwissProt ID) peptide derived from human CDK1/CDC2 around the phosphorylation site of

Thr14. AA range:1-50

产品应用 (Application)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab04432



稀释比 (Dilution Ratio) WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight) 34kDa


研究背景 (Background)

cyclin dependent kinase 1(CDK1) Homo sapiens
The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009],catalytic activity:ATP + [DNA-directed RNA polymerase] = ADP + [DNA-directed RNA polymerase] phosphate.,catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Phosphorylation at Thr-14 or Tyr-15 inactivates the enzyme, while phosphorylation at Thr-161 activates it., function: Plays a key role in the control of the eukaryotic cell cycle. It is required in higher cells for entry into S-phase and mitosis. p34 is a component of the kinase complex that phosphorylates the repetitive C-terminus of RNA polymerase II., similarity: Belongs to the protein kinase superfamily, similarity: Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family, CDC2/CDKX subfamily, similarity: Contains 1 protein kinase domain, subunit: Forms a stable but noncovalent complex with a regulatory subunit and with a cyclin. Interacts with DLGAP5. Isoform 2 is unable to complex with cyclin B1 and also fails to bind to the CDK inhibitor p21. Interacts with catalytically active CCNB1 and RALBP1 during mitosis to form an endocytotic complex during interphase.,

研究领域(Research Area)

Cell Cycle G1S;Cell Cycle G2M DNA;Oocyte meiosis;p53;Gap junction;Progesterone-mediated oocyte maturation;

图片 (Image Data)




Immunofluorescence analysis of COS7 cells, using CDK1/CDC2 (Phospho-Thr14) Antibody. The picture on the right is

E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838




Antibody

产品货号: APRab04432




Western blot analysis of lysates from HepG2 cells treated with Forskolin 40nM 30 ', using CDK1/CDC2 (Phospho-Thr14)

Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-Cdk1/2/3 (T14) Polyclonal Antibody diluted at 1: 2000



Western Blot analysis of COS7 cells using Phospho-Cdk1/2/3 (T14) Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838