Antibody

产品货号: APRab04458



产品概述 (Summary)

产品名称 (Production Name) Chk2 (phospho Thr383) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Mouse, Rat, Monkey

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CHEK2

CHEK2; CDS1; CHK2; RAD53; Serine/threonine-protein kinase Chk2; CHK2 別名 (Alternative Names)

checkpoint homolog; Cds1 homolog; Hucds1; hCds1; Checkpoint kinase 2

基因 ID (Gene ID) 11200.0

O96017.The antiserum was produced against synthesized peptide derived

蛋白 ID (SwissProt ID) from human Chk2 around the phosphorylation site of Thr383. AA range:356-

405

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

Antibody

产品货号: APRab04458



蛋白分子量 (Molecular Weight) 60kDa

研究背景 (Background)

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutaticatalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,disease:Defects in CHEK2 are associated with Li-Fraumeni syndrome 2 (LFS2) [MIM:609265]; a highly penetrant familial cancer phenotype usually associated with inherited mutations in p53/TP53, disease: Defects in CHEK2 are found in some patients with osteosarcoma (OSRC) [MIM:259500], disease: Defects in CHEK2 are found in some patients with prostate cancer (CaP) [MIM:176807], enzyme regulation: Rapidly phosphorylated on Thr-68 by MLTK in response to DNA damage and to replication block. Kinase activity is also up-regulated by autophosphorylation, function: Regulates cell cycle checkpoints and apoptosis in response to DNA damage, particularly to DNA double-strand breaks. Inhibits CDC25C phosphatase by phosphorylation on 'Ser-216', preventing the entry into mitosis. May also play a role in meiosis. Regulates the TP53 tumor suppressor through phosphorylation at 'Thr-18' and 'Ser-20', similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CHK2 subfamily, similarity: Contains 1 FHA domain, similarity: Contains 1 protein kinase domain, subcellular location: Isoform 10 is present throughout the cell, tissue specificity: High expression is found in testis, spleen, colon and peripheral blood leukocytes. Low expression is found in other tissues.,

研究领域 (Research Area)

Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;p53;


图片 (Image Data)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838


Antibody

产品货号: APRab04458






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using Chk2 (Phospho-Thr383) Antibody



Immunofluorescence analysis of HeLa cells, using Chk2 (Phospho-Thr383) Antibody. The picture on the right is blocked with the phospho peptide.




Western blot analysis of lysates from COS7 cells treated with UV 30 ', using Chk2 (Phospho-Thr383) Antibody. The lane on the right is blocked with the phospho peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

Antibody

产品货号: APRab04458





Western Blot analysis of Hela cells using Phospho-Chk2 (T383) Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838