

产品名称: c-Myc (phospho Thr58) Rabbit Polyclonal Antibody
产品货号: APRab04481



产品概述 (Summary)

产品名称 (Production Name)	c-Myc (phospho Thr58) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MYC
别名 (Alternative Names)	MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64
基因 ID (Gene ID)	4609.0
蛋白 ID (SwissProt ID)	P01106.The antiserum was produced against synthesized peptide derived from human Myc around the phosphorylation site of Thr58. AA range:25-74

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000,IP 1:20-1:50
----------------------	---

产品名称: c-Myc (phospho Thr58) Rabbit Polyclonal Antibody
产品货号: APRab04481



蛋白分子量 (Molecular Weight) 50,(also ~60kDa in some samples)


研究背景 (Background)

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq, Jul 2008],disease:A chromosomal aberration involving MYC may be a cause of a form of B-cell chronic lymphocytic leukemia. Translocation t(8;12)(q24;q22) with BTG1.,disease:Overexpression of MYC is implicated in the etiology of a variety of hematopoietic tumors.,function:Participates in the regulation of gene transcription. Binds DNA both in a non-specific manner and also specifically to recognizes the core sequence 5'-CAC[GA]TG-3'. Seems to activate the transcription of growth-related genes.,online information:Myc entry,PTM:Phosphorylated by PRKDC.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Binds DNA as a heterodimer with MAX. Interacts with TAF1C and SPAG9. Interacts with PARP10. Interacts with KDM5A and KDM5B.,

研究领域 (Research Area)


Stem cell pathway; Cell_Cycle_G1S;Cell_Cycle_G2M_DNA; WNT;WNT-T CELL;β-Catenin; ErbB/HER; MAPK_ERK_Growth;MAPK_G_Protein; PI3K/Akt; Protein_Acetylation

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Myc (Phospho-Thr58) Antibody.
The picture on the right is blocked with the phospho peptide.


产品名称: c-Myc (phospho Thr58) Rabbit Polyclonal Antibody
产品货号: APRab04481



Western blot analysis of lysates from ovary cancer, using Myc (Phospho-Thr58) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-c-Myc (T58) Polyclonal Antibody diluted at 1: 500



Western Blot analysis of 293 cells using Phospho-c-Myc (T58) Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .