

产品货号: APRab04518



产品概述 (Summary)

Cyclin B1 (phospho Ser147) Rabbit Polyclonal Antibody 产品名称 (Production Name)

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA 种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

Unconjugated 偶联物 (Conjugation) 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CCNB1

别名 (Alternative Names) CCNB1; CCNB; G2/mitotic-specific cyclin-B1

基因 ID (Gene ID) 891.0

P14635.The antiserum was produced against synthesized peptide derived

from human Cyclin B1 around the phosphorylation site of Ser147. AA 蛋白 ID (SwissProt ID)

range:121-170

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 60kDa

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838



产品货号: APRab04518




研究背景 (Background)

The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq, Jul 2008], developmental stage:Accumulates steadily during G2 and is abruptly destroyed at mitosis, function: Essential for the control of the cell cycle at the G2/M (mitosis) transition.,PTM:Ubiquitinated by the SCF(NIPA) complex during interphase, leading to its destruction. Not ubiquitinated during G2/M phases, similarity: Belongs to the cyclin family, similarity: Belongs to the cyclin family. Cyclin AB subfamily, subunit: Interacts with the CDC2 protein kinase to form a serine/threonine kinase holoenzyme complex also known as maturation promoting factor (MPF). The cyclin subunit imparts substrate specificity to the complex. Binds HEI10. Interacts with catalytically active RALBP1 and CDC2 during mitosis to form an endocytotic complex during interphase.,

研究领域(Research Area)

AMPK

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Cyclin B1 (Phospho-Ser147) Antibody


E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838






产品货号: APRab04518





Immunohistochemistry analysis of paraffin-embedded human placenta, using Cyclin B1 (Phospho-Ser147) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells treated with UV 15 ', using Cyclin B1 (Phospho-Ser147) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838