

产品名称: Dematin (phospho Ser403) Rabbit Polyclonal Antibody
产品货号: APRab04546



产品概述 (Summary)

产品名称 (Production Name)	Dematin (phospho Ser403) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	EPB49
别名 (Alternative Names)	EPB49; DMT; Dematin; Erythrocyte membrane protein band 4.9
基因 ID (Gene ID)	2039.0
蛋白 ID (SwissProt ID)	Q08495.The antiserum was produced against synthesized peptide derived from human Dematin around the phosphorylation site of Ser403. AA range:356-405

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	55kDa

产品名称: Dematin (phospho Ser403) Rabbit Polyclonal Antibody
产品货号: APRab04546




研究背景 (Background)

The protein encoded by this gene is an actin binding and bundling protein that plays a structural role in erythrocytes, by stabilizing and attaching the spectrin/actin cytoskeleton to the erythrocyte membrane in a phosphorylation-dependent manner. This protein contains a core domain in the N-terminus, and a headpiece domain in the C-terminus that binds F-actin. When purified from erythrocytes, this protein exists as a trimer composed of two 48 kDa polypeptides and a 52 kDa polypeptide. The different subunits arise from alternative splicing in the 3' coding region, where the headpiece domain is located. Disruption of this gene has been correlated with the autosomal dominant Marie Unna hereditary hypotrichosis disease, while loss of heterozygosity of this gene is thought to play a role in prostate cancer progression. Alternative splicing results in multiple transcript variants encoding didomain: Consists of a large core fragment, the amino-terminal portion, and a small headpiece, the C-terminal portion. The headpiece can bind but cannot bundle actin filaments., domain: Contains at least two actin-binding sites, one in the headpiece domain and one in the amino-terminal portion., function: Actin-bundling protein. May function in mitogen-activated protein kinase pathway., PTM: Actin-bundling activity is abolished upon phosphorylation by cAMP-dependent protein kinase., PTM: The N-terminus is blocked., similarity: Belongs to the villin/gelsolin family., similarity: Contains 1 HP (headpiece) domain., subunit: Exists in solution as a trimer of two short isoforms and one long isoform linked by disulfide bonds (Probable). Interacts with RASGRF2., tissue specificity: Heart, brain, lung, skeletal muscle, and kidney.,


研究领域 (Research Area)

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human heart, using Dematin (Phospho-Ser403) Antibody. The picture on the right is blocked with the phospho peptide.

产品名称: Dematin (phospho Ser403) Rabbit Polyclonal Antibody
产品货号: APRab04546



Western blot analysis of lysates from HepG2 cells treated with Insulin 0.01U/ml 15 ' ; using Dematin (Phospho-Ser403) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .