

产品名称: FoxO4 (phospho Thr451) Rabbit Polyclonal Antibody
产品货号: APRab04706



产品概述 (Summary)

产品名称 (Production Name)	FoxO4 (phospho Thr451) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FOXO4
别名 (Alternative Names)	FOXO4; AFX; AFX1; MLLT7; Forkhead box protein O4; Fork head domain transcription factor AFX1
基因 ID (Gene ID)	4303.0
蛋白 ID (SwissProt ID)	P98177.The antiserum was produced against synthesized peptide derived from human FOXO4 around the phosphorylation site of Thr451. AA range:417-466

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:20000-1:40000
----------------------	---------------------------------------

产品名称: FoxO4 (phospho Thr451) Rabbit Polyclonal Antibody
产品货号: APRab04706



蛋白分子量 (Molecular Weight) 55kDa


研究背景 (Background)

This gene encodes a member of the O class of winged helix/forkhead transcription factor family. Proteins encoded by this class are regulated by factors involved in growth and differentiation indicating they play a role in these processes. A translocation involving this gene on chromosome X and the homolog of the Drosophila trithorax gene, encoding a DNA binding protein, located on chromosome 11 is associated with leukemia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2010],disease:A chromosomal aberration involving FOXO4 is found in acute leukemias. Translocation t(X;11)(q13;q23) with MLL/HRX. The result is a rogue activator protein.,function:Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGFBP1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle.,pharmaceutical:A constitutively active FOXO4 mutant where phosphorylation sites Thr-32, Ser-187 and Ser-262 have been mutated to alanine may have therapeutic potential in ERBB2/HER2-overexpressing cancers as it inhibits ERBB2-mediated cell survival, transformation and tumorigenicity.,PTM:Acetylation by CBP, which is induced by peroxidase stress, inhibits transcriptional activity. Deacetylation by SIRT1 is NAD-dependent and stimulates transcriptional activity.,PTM:Phosphorylation by PKB/AKT1 inhibits transcriptional activity and is responsible for cytoplasmic localization.,similarity:Contains 1 fork-head DNA-binding domain.,subcellular location:When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation.,subunit:Interacts with CBP, MYOCD, SIRT1, SRF and YWHAZ. Acetylated by CBP and deacetylated by SIRT1. Binding of YWHAZ inhibits DNA-binding.,tissue specificity:Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform zeta is most abundant in the liver, kidney, and pancreas.,

研究领域 (Research Area)

Insulin Receptor; B Cell Receptor; Protein_Acetylation


图片 (Image Data)



产品名称: FoxO4 (phospho Thr451) Rabbit Polyclonal Antibody
产品货号: AP Rab04706



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using FOXO4 (Phospho-Thr451) Antibody



Western blot analysis of lysates from HUVEC cells treated with EGF 200ng/ml 5', using FOXO4 (Phospho-Thr451) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.