

产品名称: GATA-4 (phospho Ser105) Rabbit Polyclonal Antibody
产品货号: APRab04727



产品概述 (Summary)

产品名称 (Production Name)	GATA-4 (phospho Ser105) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GATA4
别名 (Alternative Names)	GATA4; Transcription factor GATA-4; GATA-binding factor 4
基因 ID (Gene ID)	2626.0
蛋白 ID (SwissProt ID)	P43694. The antiserum was produced against synthesized peptide derived from human GATA4 around the phosphorylation site of Ser105. AA range: 71-120

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:200-1:1000, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: GATA-4 (phospho Ser105) Rabbit Polyclonal Antibody
产品货号: APRab04727




研究背景 (Background)

This gene encodes a member of the GATA family of zinc-finger transcription factors. Members of this family recognize the GATA motif which is present in the promoters of many genes. This protein is thought to regulate genes involved in embryogenesis and in myocardial differentiation and function, and is necessary for normal testicular development. Mutations in this gene have been associated with cardiac septal defects. Additionally, alterations in gene expression have been associated with several cancer types. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015],disease:Defects in GATA4 are the cause of atrial septal defect type 2 (ASD2) [MIM:607941]. ASD2 is a congenital heart malformation characterized by incomplete closure of the wall between the atria resulting in blood flow from the left to the right atria. ASD2 patients show other heart abnormalities including ventricular and atrioventricular septal defects, pulmonary valve thickening or insufficiency of the cardiac valves. ASD2 is not associated with defects in the cardiac conduction system or non-cardiac abnormalities.,function:Transcriptional activator. Binds to the consensus sequence 5'-AGATAG-3'. Acts as a transcriptional activator of ANF in cooperation with NKX2-5.,similarity:Contains 2 GATA-type zinc fingers.,subunit:Interacts with ZNF260 (By similarity). Interacts with the homeobox domain of NKX2-5 through its C-terminal zinc finger. Also interacts with JARID2 which represses its ability to activate transcription of ANF. Interacts with NFATC4.,


研究领域 (Research Area)

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GATA4 (Phospho-Ser105) Antibody

产品名称: GATA-4 (phospho Ser105) Rabbit Polyclonal Antibody
产品货号: APRab04727



Immunofluorescence analysis of HepG2 cells, using GATA4 (Phospho-Ser105) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .