

产品名称: HDAC8 (phospho Ser39) Rabbit Polyclonal Antibody
产品货号: APRab04768



产品概述 (Summary)

产品名称 (Production Name)	HDAC8 (phospho Ser39) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	HDAC8
别名 (Alternative Names)	HDAC8; HDACL1; CDA07; Histone deacetylase 8; HD8
基因 ID (Gene ID)	55869.0
蛋白 ID (SwissProt ID)	Q9BY41.The antiserum was produced against synthesized peptide derived from human HDAC8 around the phosphorylation site of Ser39. AA range:5-54

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: HDAC8 (phospho Ser39) Rabbit Polyclonal Antibody
产品货号: APRab04768




研究背景 (Background)

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009],catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.,caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.,miscellaneous:Its activity is inhibited by trichostatin A (TSA) and butyrate, two well known histone deacetylase inhibitors.,similarity:Belongs to the histone deacetylase family. Type 1 subfamily.,subcellular location:Excluded from the nucleoli.,subunit:Interacts with PEPB2-MYH11, a fusion protein consisting of the 165 N-terminal residues of CBF-beta (PEPB2) with the tail region of MYH11 produced by the inversion Inv(16)(p13q22), a translocation associated with acute myeloid leukemia of M4EO subtype. The PEPB2-MYH1 fusion protein also interacts with RUNX1, a well known transcriptional regulator, suggesting that the interaction with HDAC8 may participate in the conversion of RUNX1 into a constitutive transcriptional repressor. Interacts with CBFA2T3.,tissue specificity:Weakly expressed in most tissues. Expressed at higher level in heart, brain, kidney and pancreas.,

研究领域 (Research Area)


Protein_Acetylation

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using HDAC8 (Phospho-Ser39) Antibody.
The picture on the right is blocked with the phospho peptide.


产品名称: HDAC8 (phospho Ser39) Rabbit Polyclonal Antibody
产品货号: APRab04768



Western blot analysis of lysates from NIH/3T3 cells, using HDAC8 (Phospho-Ser39) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-HDAC8 (S39) Polyclonal Antibody diluted at 1: 500



Western Blot analysis of VEC cells using Phospho-HDAC8 (S39) Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .