

产品名称: HP1 α (phospho Ser92) Rabbit Polyclonal Antibody
产品货号: APRab04791



产品概述 (Summary)

产品名称 (Production Name)	HP1 α (phospho Ser92) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CBX5
别名 (Alternative Names)	CBX5; HP1A; Chromobox protein homolog 5; Antigen p25; Heterochromatin protein 1 homolog alpha; HP1 alpha
基因 ID (Gene ID)	23468.0
蛋白 ID (SwissProt ID)	P45973. The antiserum was produced against synthesized peptide derived from human HP1 alpha around the phosphorylation site of Ser92. AA range: 58-107

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	

产品名称: HP1 α (phospho Ser92) Rabbit Polyclonal Antibody
产品货号: APRab04791




研究背景 (Background)

This gene encodes a highly conserved nonhistone protein, which is a member of the heterochromatin protein family. The protein is enriched in the heterochromatin and associated with centromeres. The protein has a single N-terminal chromodomain which can bind to histone proteins via methylated lysine residues, and a C-terminal chromoshadow-domain (CSD) which is responsible for the homodimerization and interaction with a number of chromatin-associated nonhistone proteins. The encoded product is involved in the formation of functional kinetochore through interaction with essential kinetochore proteins. The gene has a pseudogene located on chromosome 3. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul 2008], function: Component of heterochromatin. Recognizes and binds histone H3 tails methylated at 'Lys-9', leading to epigenetic repression. Can interact with lamin B receptor (LBR). This interaction can contribute to the association of the heterochromatin with the inner nuclear membrane. Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins., PTM: Phosphorylation of HP1 and LBR may be responsible for some of the alterations in chromatin organization and nuclear structure which occur at various times during the cell cycle (By similarity). Phosphorylated during interphase and possibly hyper-phosphorylated during mitosis., similarity: Contains 2 chromo domains., subcellular location: Component of centromeric and pericentromeric heterochromatin. Associates with chromosomes during mitosis. Associates specifically with chromatin during metaphase and anaphase., subunit: Interacts with SUV420H1 and SUV420H2 (By similarity). Interacts directly with ATRX, CHAF1A, LBR, NIPBL, SP100, STAM2 and TRIM28 via the chromoshadow domain. Can interact directly with CBX3 via the chromoshadow domain. Interacts with histone H3 methylated at 'Lys-9'. Interacts with MIS12 and C20orf127. Interacts with HP1BP3.,

研究领域 (Research Area)


Epigenetics and Nuclear Signaling

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HP1 alpha (Phospho-Ser92) Antibody

产品名称: HP1 α (phospho Ser92) Rabbit Polyclonal Antibody
产品货号: APRab04791



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using HP1 alpha (Phospho-Ser92) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .