

**产品名称: IKK $\beta$  (phospho Tyr199) Rabbit Polyclonal Antibody**  
**产品货号: APRab04829**



## 产品概述 (Summary)

|                        |                                                         |
|------------------------|---------------------------------------------------------|
| 产品名称 (Production Name) | IKK $\beta$ (phospho Tyr199) Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody                              |
| 宿主 (Host)              | Rabbit                                                  |
| 应用 (Application)       | WB,IHC,ICC/IF,ELISA                                     |
| 种属反应性 (Reactivity)     | Human,Mouse,Rat                                         |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Phosphorylated                                                                                    |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                                                                                                                         |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | IKBKB                                                                                                                                                                                                   |
| 别名 (Alternative Names) | IKBKB; IKKB; Inhibitor of nuclear factor kappa-B kinase subunit beta; I-kappa-B-kinase beta; IKK-B; IKK-beta; IkbKB; I-kappa-B kinase 2; IKK2; Nuclear factor NF-kappa-B inhibitor kinase beta; NFKB1KB |
| 基因 ID (Gene ID)        | 3551.0                                                                                                                                                                                                  |
| 蛋白 ID (SwissProt ID)   | O14920.The antiserum was produced against synthesized peptide derived from human IKK-beta around the phosphorylation site of Tyr199. AA range:166-215                                                   |

## 产品应用 (Application)

**产品名称: IKK $\beta$  (phospho Tyr199) Rabbit Polyclonal Antibody**  
**产品货号: APRab04829**

---

|                                 |                                                                        |
|---------------------------------|------------------------------------------------------------------------|
| <b>稀释比 (Dilution Ratio)</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000 |
| <b>蛋白分子量 (Molecular Weight)</b> | 85kDa                                                                  |

## 研究背景 (Background)


The protein encoded by this gene phosphorylates the inhibitor in the inhibitor/NF-kappa-B complex, causing dissociation of the inhibitor and activation of NF-kappa-B. The encoded protein itself is found in a complex of proteins. Several transcript variants, some protein-coding and some not, have been found for this gene. [provided by RefSeq, Sep 2011],catalytic activity:ATP + [I-kappa-B protein] = ADP + [I-kappa-B phosphoprotein],function:Acts as part of the IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. Also phosphorylates NCOA3.,PTM:Ubiquitination on 'Ser-163' modulates phosphorylation on C-terminal serine residues.,PTM:Upon cytokine stimulation, phosphorylated on Ser-177 and Ser-181 by MEKK1 and/or MAP3K14/NIK; which enhances activity. Once activated, autophosphorylates on the C-terminal serine cluster; which decreases activity and prevents prolonged activation of the inflammatory response.,PTM:Yersinia yopJ may acetylate Ser/Thr residues, preventing phosphorylation and activation, which blocks the I-kappa-B signaling pathway.,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. I-kappa-B kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Component of the I-kappa-B-kinase (IKK) core complex consisting of CHUK, IKBKB and IKBKG; probably four alpha/CHUK-beta/IKBKB dimers are associated with four gamma/IKBKG subunits. The IKK core complex seems to associate with regulatory or adapter proteins to form a IKK-signalosome holo-complex. Part of a complex composed of NCOA2, NCOA3, CHUK/IKKA, IKBKB, IKBKG and CREBBP. Part of a 70-90 kDa complex at least consisting of CHUK/IKKA, IKBKB, NFKBIA, RELA, IKBKAP and MAP3K14. Interacts with SQSTM1 through PRKCZ or PRKCI. Forms an NGF-induced complex with IKBKB, PRKCI and TRAF6. May interact with MAVS/IPS1. Interacts with NALP2. Interacts with TICAM1. Interacts with Yersinia yopJ. Interacts with FAF1; the interaction disrupts the IKK complex formation. Interacts with ATM. Part of a ternary complex consisting of TANK, IKBKB and IKBKG. Interacts with NIBP; the interaction is direct.,tissue specificity:Highly expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis and peripheral blood.,

## 研究领域 (Research Area)


MAPK\_ERK\_Growth;MAPK\_G\_Protein;Chemokine;Apoptosis\_Inhibition;Apoptosis\_Mitochondrial;Apoptosis\_Overview;Toll\_Like;NOD-like receptor;RIG-I-like receptor;Cytosolic DNA-sensing pathway;T\_Cell\_Receptor;B\_Cell\_Antigen;Neurotrophin;Insulin\_Receptor;Adipocytokine;Type II diabetes mellitus;Epithelial cell signaling in Helicobacter pylori infection;Pathways in cancer;Pancreatic cancer;Prostate cancer;Chronic myeloid leukemia;Acute myeloid leukemia;Small cell lung cancer;

## 图片 (Image Data)


产品名称: IKK $\beta$  (phospho Tyr199) Rabbit Polyclonal Antibody  
产品货号: APRab04829



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IKK-beta (Phospho-Tyr199) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using IKK-beta (Phospho-Tyr199) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with TNF-a 20ng/ml + Calyculin A 50nM 5', using IKK-beta (Phospho-Tyr199) Antibody. The lane on the right is blocked with the phospho peptide.

## 注意事项 (Note)

For research use only.