

产品名称: IRAK-1 (phospho Ser376) Rabbit Polyclonal Antibody
产品货号: APRab04864



产品概述 (Summary)

产品名称 (Production Name)	IRAK-1 (phospho Ser376) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IRAK1
别名 (Alternative Names)	IRAK1; IRAK; Interleukin-1 receptor-associated kinase 1; IRAK-1
基因 ID (Gene ID)	3654.0
蛋白 ID (SwissProt ID)	P51617. The antiserum was produced against synthesized peptide derived from human IRAK1 around the phosphorylation site of Ser376. AA range: 342-391

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	

产品名称: IRAK-1 (phospho Ser376) Rabbit Polyclonal Antibody
产品货号: APRab04864




研究背景 (Background)

This gene encodes the interleukin-1 receptor-associated kinase 1, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. This gene is partially responsible for IL1-induced upregulation of the transcription factor NF-kappa B. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., function: Binds to the IL-1 type I receptor following IL-1 engagement, triggering intracellular signaling cascades leading to transcriptional up-regulation and mRNA stabilization. Isoform 1 binds rapidly but is then degraded allowing isoform 2 to mediate a slower, more sustained response to the cytokine. Isoform 2 is inactive suggesting that the kinase activity of this enzyme is not required for IL-1 signaling. Once phosphorylated, IRAK1 recruits the adapter protein PELI1., PTM: Autophosphorylated or is transphosphorylated by IRAK4 following recruitment to the IL-1RI. In the case of isoform 1, this is linked to ubiquitination and degradation., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Pelle subfamily., similarity: Contains 1 protein kinase domain., subunit: IL-1 stimulation leads to the formation of a signaling complex which dissociates from the IL-1 receptor following the binding of PELI1. Interacts with IL1RL1. Interacts with IRAK1BP1., tissue specificity: Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.,

研究领域 (Research Area)

Apoptosis_Inhibition; Apoptosis_Mitochondrial; Apoptosis_Overview; Toll_Like; Neurotrophin;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain, using IRAK1 (Phospho-Ser376) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .

产品名称: IRAK-1 (phospho Ser376) Rabbit Polyclonal
Antibody
产品货号: APRab04864

