产品名称: IRAK-1 (phospho Thr387) Rabbit Polyclonal

Antibody

产品货号: APRab04867



#### 产品概述 (Summary)

产品名称 (Production Name) IRAK-1 (phospho Thr387) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,IHC

种属反应性 (Reactivity) Human, Mouse, Rat

### 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

### 免疫原信息 (Immunogen)

基因名 (Gene Name) IRAK1

別名 (Alternative Names) IRAK1; IRAK; Interleukin-1 receptor-associated kinase 1; IRAK-1

基因 ID (Gene ID) 3654.0

P51617.The antiserum was produced against synthesized peptide derived

蛋白 ID (SwissProt ID) from human IRAK1 around the phosphorylation site of Thr387. AA range:353-

402

# 产品应用(Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:50-1:300

蛋白分子量(Molecular Weight)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: IRAK-1 (phospho Thr387) Rabbit Polyclonal

**Antibody** 

产品货号: APRab04867




#### 研究背景 (Background)

This gene encodes the interleukin-1 receptor-associated kinase 1, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. This gene is partially responsible for IL1-induced upregulation of the transcription factor NF-kappa B. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, function:Binds to the IL-1 type I receptor following IL-1 engagement, triggering intracellular signaling cascades leading to transcriptional up-regulation and mRNA stabilization. Isoform 1 binds rapidly but is then degraded allowing isoform 2 to mediate a slower, more sustained response to the cytokine. Isoform 2 is inactive suggesting that the kinase activity of this enzyme is not required for IL-1 signaling. Once phosphorylated, IRAK1 recruits the adapter protein PELI1,PTM:Autophosphorylated or is transphosphorylated by IRAK4 following recruitment to the IL-1RI. In the case of isoform 1, this is linked to ubiquitination and degradation, similarity:Belongs to the protein kinase superfamily, similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Pelle subfamily, similarity:Contains 1 protein kinase domain, subunit:IL-1 stimulation leads to the formation of a signaling complex which dissociates from the IL-1 receptor following the binding of PELI1. Interacts with IL1RL1. Interacts with IRAK1BP1, tissue specificity:Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.

### 研究领域 (Research Area)

Apoptosis Inhibition; Apoptosis Mitochondrial; Apoptosis Overview; Toll Like; Neurotrophin;

#### 图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IRAK1 (Phospho-Thr387) Antibody

# 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: IRAK-1 (phospho Thr387) Rabbit Polyclonal



Antibody

产品货号: APRab04867

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838