

产品货号: APRab04970

#### 产品概述 (Summary)

产品名称 (Production Name) MAPKAPK-2 (phospho Ser272) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse,Rat

### 产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) MAPKAPK2

MAPKAPK2; MAP kinase-activated protein kinase 2; MAPK-activated protein 别名 (Alternative Names)

kinase 2; MAPKAP kinase 2; MAPKAP-K2; MAPKAPK-2; MK-2; MK2

基因 ID (Gene ID) 9261.0

P49137.The antiserum was produced against synthesized peptide derived

**蛋白ID (SwissProt ID)** from human MAPKAPK2 around the phosphorylation site of Ser272. AA

range:238-287

# 产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: MAPKAPK-2 (phospho Ser272) Rabbit Polyclonal

**Antibody** 

产品货号: APRab04970



蛋白分子量 (Molecular Weight) 45-52kDa


### 研究背景 (Background)

This gene encodes a member of the Ser/Thr protein kinase family. This kinase is regulated through direct phosphorylation by p38 MAP kinase. In conjunction with p38 MAP kinase, this kinase is known to be involved in many cellular processes including stress and inflammatory responses, nuclear export, gene expression regulation and cell proliferation. Heat shock protein HSP27 was shown to be one of the substrates of this kinase in vivo. Two transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein,,enzyme regulation:Seems to be activated by two distinct pathways: the first involves the stimulation of p42/p44 MAPK by growth factors, the second, triggered by stress and heat shock, depends on the activation of MPK2 and upstream MAPKK/MAPKKK,function:Its physiological substrate seems to be the small heat shock protein (HSP27/HSP25). In vitro can phosphorylate glycogen synthase at 'Ser-7' and tyrosine hydroxylase (on 'Ser-19' and 'Ser-40'). This kinase phosphorylates Ser in the peptide sequence, Hyd-X-R-X(2)-S, where Hyd is a large hydrophobic residue (By similarity). Mediates both ERK and p38 MAPK/MAPK14 dependent neutrophil responses. Participates in TNF alpha-stimulated exocytosis of secretory vesicles in neutrophils. Plays a role in phagocytosis-induced respiratory burst activity.,PTM:Phosphorylated and activated by MAP kinase, similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family, similarity:Contains 1 protein kinase domain., subunit:Interacts with PHC2, tissue specificity:Expressed in all tissues examined.,

#### 研究领域(Research Area)

MAPK ERK Growth; MAPK G Protein; VEGF; Neurotrophin;

### 图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human brain, using MAPKAPK2 (Phospho-Ser272) Antibody. The picture on the right is blocked with the phospho peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838



产品货号: APRab04970



Western blot analysis of lysates from RAW264.7 cells treated with UV 15 ', using MAPKAPK2 (Phospho-Ser272) Antibody.

The lane on the right is blocked with the phospho peptide.

#### 注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838