

产品名称: Mnk1 (phospho Thr250) Rabbit Polyclonal Antibody
产品货号: APRab05030



产品概述 (Summary)

产品名称 (Production Name)	Mnk1 (phospho Thr250) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MKNK1
别名 (Alternative Names)	MKNK1; MNK1; MAP kinase-interacting serine/threonine-protein kinase 1; MAP kinase signal-integrating kinase 1; MAPK signal-integrating kinase 1; Mnk1
基因 ID (Gene ID)	8569.0
蛋白 ID (SwissProt ID)	Q9BUB5. The antiserum was produced against synthesized peptide derived from human MNK1 around the phosphorylation site of Thr250. AA range: 216-265

产品应用 (Application)

产品名称: Mnk1 (phospho Thr250) Rabbit Polyclonal Antibody
产品货号: APRab05030



稀释比 (Dilution Ratio)

IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:10000-1:20000

蛋白分子量 (Molecular Weight)


研究背景 (Background)

MAP kinase interacting serine/threonine kinase 1(MKNK1) Homo sapiens This gene encodes a Ser/Thr protein kinase that interacts with, and is activated by ERK1 and p38 mitogen-activated protein kinases, and thus may play a role in the response to environmental stress and cytokines. This kinase may also regulate transcription by phosphorylating eIF4E via interaction with the C-terminal region of eIF4G. Alternatively spliced transcript variants have been noted for this gene. [provided by RefSeq, Jan 2012], catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., enzyme regulation: Phosphorylated and activated by the p38 kinases and kinases in the Erk pathway., function: May play a role in the response to environmental stress and cytokines. Appears to regulate transcription by phosphorylating EIF4E, thus increasing the affinity of this protein for the 7-methylguanosine-containing mRNA cap., PTM: Dual phosphorylation of Thr-250 and Thr-255 activates the kinase. Phosphorylation of Thr-385 activates the kinase., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family., similarity: Contains 1 protein kinase domain., subunit: Interacts with the C-terminal regions of EIF4G1 and EIF4G2. Also binds to dephosphorylated ERK1 and ERK2, and to the p38 kinases., tissue specificity: Ubiquitous.,

研究领域 (Research Area)

MAPK_ERK_Growth; MAPK_G_Protein; Insulin_Receptor;

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MNK1 (Phospho-Thr250) Antibody

注意事项 (Note)

For research use only .